

Name: _____

Date: _____

Score: _____

Vaccine Administration Post Test

Each question is worth 10 points. Minimum passing score: 80%

1. What are the routes of administration for routine childhood and adult vaccination?
 - A. Intramuscular, subcutaneous, and intradermal.
 - B. Subcutaneous, intravenous, and intramuscular.
 - C. Intramuscular, intranasal, oral, and subcutaneous.
2. True or False: The Vastus Lateralis is the correct site for IM injections for infants.
3. True or False: LAIV for influenza vaccination can only be administered to healthy, non-pregnant persons aged 2 through 49 years of age.
4. What factors should be considered when determining the needle size and site for an intramuscular injection?
 - A. Age.
 - B. Size of muscle.
 - C. Vaccine volume.
 - D. All of the above.
5. True or False: The patient must sniff during vaccine administration of LAIV.
6. True or False: It is necessary to aspirate before injecting a vaccine.
7. What are the two vaccines that may be given IM and SC?
 - A. Influenza and DTaP
 - B. Pneumococcal polysaccharide and IPV.
 - C. Hib and MMR
8. Which site is never recommended for immunizations?
 - A. Deltoid.
 - B. Anterolateral Thigh.
 - C. Vastus Lateralis.
 - D. Gluteus Maximus.

9. True or False: Syringes pre-filled by staff members should be used on the same day they are filled.

10. True or False: When multiple injections are administered, you should use the same site for each one.