

## Vaccine Administration Post Test Answer Key

Each question is worth 10 points. Minimum passing score: 80%

1. What are the routes of administration for routine childhood and adult vaccination?
  - A. Intramuscular, subcutaneous, and intradermal.
  - B. Subcutaneous, intravenous, and intramuscular.
  - C. Intramuscular, intranasal, oral, Intradermal and subcutaneous.
2. True or False: The Vastus Lateralis is the correct site for IM injections for newborns and infants.
3. True or False: LAIV can only be administered to healthy, non-pregnant persons aged 2 through 49 years of age.
4. What factors should be considered when determining the needle size and site for an intramuscular injection?
  - A. Age.
  - B. Size of muscle.
  - C. Vaccine volume.
  - D. All of the above.
5. True or False: The patient must sniff during vaccine administration of LAIV.
6. True or False: It is necessary to aspirate before injecting a vaccine.
7. Which vaccines may be given IM and SC?
  - A. Influenza and DTaP
  - B. Pneumococcal polysaccharide, and IPV.
  - C. Hib and MMR
8. Which site is never recommended for immunizations?
  - A. Deltoid.
  - B. Anterolateral Thigh.
  - C. Vastus Lateralis.
  - D. Gluteus Maximus.
9. True or False: Syringes pre-filled by staff members should be used on the same day they are filled.
10. True or False: When multiple injections are administered, you should use the same site for each one.