

2024

HIV SURVEILLANCE SUMMARY REPORT, GEORGIA

HIV Epidemiology Unit



The **HIV Surveillance Summary, Georgia 2024** is published by the Georgia Department of Public Health (DPH), Division of Epidemiology, Viral Hepatitis HIV STI Epidemiology Section, HIV Epidemiology Unit, 200 Piedmont Avenue SE, West Tower Floor 10, Atlanta, GA 30334. Data are presented from known diagnoses and laboratory reports entered into the Georgia Enhanced HIV/AIDS Reporting System (eHARS). All data are provisional.

The *HIV Surveillance Summary, Georgia 2024* is not copyrighted and may be used and reproduced without permission. Citation of the source is, however, appreciated.

SUGGESTED CITATION: Georgia Department of Public Health, HIV Epidemiology Unit, *2024 Georgia HIV Surveillance Summary*, <https://dph.georgia.gov/epidemiology/hiv-epidemiology-unit/hiv-case-surveillance-and-care-continuum> , *Published April 2026, [Accessed: date]*

ACKNOWLEDGEMENTS: Publication of this report was made possible with the contributions of the Georgia DPH HIV Epidemiology Unit’s Core HIV Surveillance staff, HIV Case Report Forms submitted by Georgia healthcare facility staff, HIV infection-related laboratory test results transmitted by laboratory facilities in Georgia, data matches with other DPH programs, and the ongoing efforts of multiple individuals from public and private sector organizations dedicated to improving surveillance, prevention, testing, and care of people with HIV infection.

This report was prepared by the following staff from the Georgia Department of Public Health: Jose Adame, MPH; Nellie Garlow, MPH; Jenna Gettings, DVM, MPH; Cherie Drenzek, DVM, MS.

Please direct all inquiries to:

Jenna Gettings, DVM, MPH
Director
Viral Hepatitis, HIV, and STI Epidemiology Section (VHHSES)
jenna.gettings@dph.ga.gov

Georgia HIV Core Surveillance Team contributors: Lauren Barrineau-Vejjajiva, Kimberly Booker, Kiswana Branch, Thelma Fannin, LaTosha Johnson, LaToya Lemons, Kim Norris, Rama Namballa, Doris Pearson, Courtney Philpot, A. Eugene Pennisi, Sylve Jones, Akilah Spratling, Erica Williams, and Gerrianna Williams.

Table of Contents

INTRODUCTION	5
TECHNICAL NOTES	6
DEFINITIONS	9
REFERENCES	10
HIV DIAGNOSES AND PEOPLE WITH HIV, GEORGIA.....	12
<i>Table 1. HIV Diagnoses, Late Diagnoses, and Stage 3 (AIDS) Diagnoses, GA, 2024</i>	12
<i>Table 2. HIV Diagnoses, Late Diagnoses, and Stage 3 (AIDS) Diagnoses among Cisgender Men, GA, 2024</i>	13
<i>Table 3. HIV Diagnoses, Late Diagnoses, and Stage 3 (AIDS) Diagnoses among Cisgender Women, GA, 2024</i>	14
<i>Table 4. HIV Diagnoses, Late Diagnoses, and Stage 3 (AIDS) Diagnoses among Transgender Persons and Persons with an Additional Gender Identity, GA, 2024</i>	15
<i>Table 5. HIV Diagnoses, Late Diagnoses, and Stage 3 (AIDS) Diagnoses, among Black or African American, non-Hispanic, Individuals, GA, 2024</i>	16
<i>Table 6. HIV Diagnoses, Late Diagnoses, and Stage 3 (AIDS) Diagnoses, among White, non-Hispanic, Individuals, GA, 2024</i>	17
<i>Table 7. HIV Diagnoses, Late Diagnoses, and Stage 3 (AIDS) Diagnoses, among Hispanic or Latino (of Any Race) Individuals, GA, 2024</i>	18
<i>Table 8. People with HIV (PWH), and PWH ever Stage 3 (AIDS), GA, 2024</i>	19
<i>Table 9. People with HIV (PWH) and PWH Ever Stage 3 (AIDS), Cisgender Men, GA, 2024</i>	20
<i>Table 10. People with HIV (PWH) and PWH Ever Stage 3 (AIDS), Cisgender Women, GA, 2024</i>	21
<i>Table 11. People with HIV (PWH), and PWH ever Stage 3 (AIDS), Transgender Persons or Persons with an Additional Gender Identity, GA, 2024</i>	22
<i>Table 12. People with HIV (PWH), and PWH ever Stage 3 (AIDS), among Black or African American, non-Hispanic, Individuals, GA, 2024</i>	23
<i>Table 13. People with HIV (PWH), and PWH ever Stage 3 (AIDS), Among White, non-Hispanic, Individuals, GA, 2024</i> ...	24
<i>Table 14. People with HIV (PWH), and PWH ever Stage 3 (AIDS), among Hispanic or Latino (of Any Race) Individuals, GA, 2024</i>	25
HIV DIAGNOSES AND PEOPLE WITH HIV BY PUBLIC HEALTH DISTRICT	26
<i>Table 15. HIV Diagnoses and People with HIV (PWH), District 1-1 (Rome), GA, 2024</i>	26
<i>Table 16. HIV Diagnoses and People with HIV (PWH), District 1-2 (Dalton), GA, 2024</i>	27
<i>Table 17. HIV Diagnoses and People with HIV (PWH), District 2 (Gainesville), GA, 2024</i>	28
<i>Table 18. HIV Diagnoses and People with HIV (PWH), District 3-1 (Cobb-Douglas), GA, 2024</i>	29
<i>Table 19. HIV Diagnoses and People with HIV (PWH), District 3-2 (Fulton), GA, 2024</i>	30

<i>Table 20. HIV Diagnoses and People with HIV (PWH), District 3-3 (Clayton), GA, 2024</i>	31
<i>Table 21. HIV Diagnoses and People with HIV (PWH), District 3-4 (GWR), GA, 2024</i>	32
<i>Table 22. HIV Diagnoses and People with HIV (PWH), District 3-5 (DeKalb), GA, 2024</i>	33
<i>Table 23. HIV Diagnoses and People with HIV (PWH), District 4 (LaGrange), GA, 2024</i>	34
<i>Table 24. HIV Diagnoses and People with HIV (PWH), District 5-1 (Dublin), GA, 2024</i>	35
<i>Table 25. HIV Diagnoses and People with HIV (PWH), District 5-2 (Macon), GA, 2024</i>	36
<i>Table 26. HIV Diagnoses and People with HIV (PWH), District 6 (Augusta), GA, 2024</i>	37
<i>Table 27. HIV Diagnoses and People with HIV (PWH), District 7 (Columbus), GA, 2024</i>	38
<i>Table 28. HIV Diagnoses and People with HIV (PWH), District 8-1 (Valdosta), GA, 2024</i>	39
<i>Table 29. HIV Diagnoses and People with HIV (PWH), District 8-2 (Albany), GA, 2024</i>	40
<i>Table 30. HIV Diagnoses and People with HIV (PWH), District 9-1 (Savannah), GA, 2024</i>	41
<i>Table 31. HIV Diagnoses and People with HIV (PWH), District 9-2 (Waycross), GA, 2024</i>	42
<i>Table 32. HIV Diagnoses and People with HIV (PWH), District 10 (Athens), GA, 2024</i>	43
HIV DIAGNOSES AND PEOPLE WITH HIV, ATLANTA MSA	44
<i>Table 33. HIV Diagnoses and People with HIV, Atlanta MSA¹, GA, 2024</i>	44
HIV DIAGNOSES AND LATE DIAGNOSES BY PUBLIC HEALTH DISTRICT AND ATLANTA MSA.....	45
<i>Table 34. HIV Diagnoses and Late Diagnoses by Public Health District and Atlanta MSA, GA, 2024</i>	45
PUBLIC HEALTH DISTRICT AND COUNTY COUNTS AND RATES	46
<i>Table 35. HIV Diagnoses and People with HIV (PWH) by Public Health District and County (Counts and rates per 100,000 population), GA, 2024</i>	46
DEATHS AMONG PEOPLE WITH HIV	51
<i>Table 36. Deaths (All Cause) among People with HIV (PWH), GA, 2024</i>	51
DISTRIBUTION OF GENERAL POPULATION BY RACE/ETHNICITY	52
<i>Table 37. Distribution of the General Population by Race and Ethnicity, GA, 2024</i>	52
SUPPLEMENTARY INFORMATION	53
<i>Additional Data on HIV in Georgia</i>	53
<i>Reporting</i>	53
<i>Requesting Data</i>	53

INTRODUCTION

HIV remains an important public health topic in the state of Georgia. As of the end of 2024, there were 66,896 people with HIV in Georgia, and 2,485 people were diagnosed in 2024. The number of people living with HIV continues to increase because of effective therapies now available, allowing people with HIV, who are diagnosed early and treated, to have long and healthy lives¹. Based on the Centers for Disease Control and Prevention's (CDC) 2023 HIV Surveillance Report, Georgia was ranked the 5th highest in the nation for the total number of new diagnoses of HIV infection among adults and adolescents after Texas, California, Florida, and New York, and ranked the 5th highest for the number of people with HIV infection, after California, New York, Florida, and Texas². Georgia ranked 2nd in the rate of HIV diagnosis among adults and adolescents after the District of Columbia, and 3rd in the rate of people with HIV after the District of Columbia and New York.

Monitoring the HIV epidemic in Georgia is essential for meeting the goals stated in the 2022–2025 National HIV/AIDS Strategy to: 1) prevent new HIV infections; 2) improve HIV-related health outcomes for people with HIV; 3) reduce HIV-related health disparities and inequities, and 4) achieve a more coordinated national response to the HIV epidemic among all partners and stakeholders³.

Information and reports from the other HIV Epidemiology Unit areas can be found here:

Georgia HIV Care Continuum: <https://dph.georgia.gov/epidemiology/hiv-epidemiology-unit/hiv-case-surveillance-and-care-continuum>

Georgia New HIV Diagnoses Trends: <https://dph.georgia.gov/epidemiology/hiv-epidemiology-unit/hiv-case-surveillance-and-care-continuum>

Georgia HIV Perinatal Exposure: <https://dph.georgia.gov/epidemiology/hiv-epidemiology-unit/perinatal-exposure-surveillance>

Georgia HIV Medical Monitoring Program: <https://dph.georgia.gov/epidemiology/hiv-epidemiology-unit/ga-mmp>

Georgia HIV Network Detection and Response: <https://dph.georgia.gov/epidemiology/hiv-epidemiology-unit/hiv-network-detection-and-response>

Georgia HIV Population Specific Reports: <https://dph.georgia.gov/epidemiology/hiv-epidemiology-unit/hiv-case-surveillance-and-care-continuum>

TECHNICAL NOTES

Georgia statutes and regulations (O.C.G.A. §31-12-2(b)) require healthcare providers (such as nurses, nurse practitioners, doctors, physician assistants) licensed in the state of Georgia to report all cases of HIV infection and/or Stage 3 (AIDS) to the Georgia DPH within seven days of diagnosis⁴. Laboratories licensed in the state of Georgia are required to report all HIV-related laboratory test results within 7 days. The information is used to monitor the HIV epidemic in Georgia and guide program planning and evaluation. The data presented in the accompanying tables are based on confidential case reports collected through the Georgia DPH enhanced HIV/AIDS Reporting System (eHARS).

AIDS reporting began in the early 1980s, and HIV reporting began in Georgia on December 31, 2003. Electronic transmission of laboratory test results began in 2011. Receipt of CD4 counts and viral load tests make it possible to estimate the HIV Care Continuum for people with a diagnosis of HIV infection in the state of Georgia (reports are available at: <https://dph.georgia.gov/epidemiology/hiv-epidemiology-unit/hiv-case-surveillance-and-care-continuum>)

This report includes surveillance data reported through December 31, 2025 for cases diagnosed through December 31, 2024. This report is limited to cases diagnosed through December 2024 to take into account delays in reporting, and to allow for obtaining follow up information on reported cases.

HIV diagnoses include people with a diagnosis of HIV infection, regardless of the stage of disease at diagnosis, who were residents of Georgia at the time of diagnosis. People with HIV are those with a residence in the state of Georgia during the given year, based on current information available in the Georgia HIV surveillance system, regardless of where they resided at the time of diagnosis. People residing in correctional facilities are included in this report and may inflate rates in certain geographic regions where there are large numbers of HIV-positive inmates.

Information on HIV transmission category is missing for approximately one eighth of cases. Multiple imputation, a statistical approach, is used to replace each missing transmission category with a set of plausible values that represent the uncertainty about the true, but missing, value⁵. The methods were applied to adult and adolescent, but not pediatric, cases. Multiple imputation is considered by the CDC to be the best method for redistribution of missing data in large databases. HIV transmission category is shown with missing data, and after multiple imputation. Cases missing race/ethnicity are not assigned a transmission category through multiple imputation. For more information, see the CDC's technical notes on multiple imputation:

https://stacks.cdc.gov/view/cdc/121127/cdc_121127_DS1.pdf

Prior to 2021, gender identity was determined based on sex assigned at birth and current gender identity (i.e., most recent data available). In 2021, the calculation for gender identity was adjusted by adding information from a third variable: whether the patient had ever had “transgender – female to male”, “transgender – male to female”, or “additional gender identity” listed as their current gender identity in their eHARS record. This was added to account for potential documentation issues, such as if a more recent case report form was incomplete and did not include updated information on a person’s transgender or additional gender identity status. As a result of this change, ascertainment increased for people whose gender identity was transgender or additional gender identity. For instance, in 2020, there were 683 transgender individuals living with HIV and in 2021, there were 893 transgender individuals and 39 individuals with an additional gender identity living with HIV. In 2024, CDC released new guidance on three recommended methods for calculating gender identity, encouraging jurisdictions to select one method that was most applicable for their jurisdiction. Georgia DPH adopted one of the recommended methods, which calculates gender identity by incorporating sex at birth and a summary variable that categorizes an individual as transgender or an additional gender identity if they have ever had transgender or an additional gender identity listed in their eHARS record. This calculation is very similar to the method Georgia DPH used in the prior year but now aligns with national standards.

Current gender identity is sometimes incomplete in eHARS records, so the numbers reported in these summary tables for transgender and additional gender identity are most likely an underestimate. Current gender identity fields which include the options of “transgender – female to male” and “transgender – male to female” were added to the case report form in 2007, thus prevalent counts of HIV among transgender people may be incomplete among people diagnosed prior to 2007.

People with a transmission category of “perinatal/other pediatric” include infants born in Georgia who had an HIV infection that was likely attributed to perinatal transmission, as well as infants born outside of Georgia who had an HIV infection that was likely attributed to perinatal transmission and were first diagnosed with HIV in Georgia. For more information on perinatal HIV exposure surveillance in Georgia, see the Perinatal HIV Exposure Surveillance Report at <https://dph.georgia.gov/epidemiology/hiv-epidemiology-unit/perinatal-exposure-surveillance>.

For data presented at the county, Public Health District, or Atlanta Metropolitan Statistical Area (MSA) level, all cells with values between 1 and 4 observations are censored in accordance with the Georgia Department of Public Health HIV Surveillance Security and Confidentiality Guidelines.

After the COVID-19 pandemic was declared a national and state emergency in March 2020, access to healthcare services, including HIV testing, prevention, and care-related services, became reduced or temporarily suspended. The number of people diagnosed with HIV decreased in 2020, at least in part because of decreased testing availability and changes in health care seeking patterns but returned to pre-COVID-19

levels in 2021. In 2022, HIV diagnoses were slightly higher than pre-pandemic levels, which may have indicated increased healthcare availability and healthcare seeking behaviors since the 2020 COVID-19-related disruptions. HIV diagnoses in 2022, 2023, and 2024 were comparable to pre-pandemic levels, suggesting that the impacts of the COVID-19 pandemic have stabilized.

DEFINITIONS

HIV DIAGNOSES are cases diagnosed between January 1 and December 31, 2024, with a Georgia address of residence at the time of HIV diagnosis, and who were reported to the Georgia Department of Public Health by December 31, 2025, regardless of stage at diagnosis. This group includes people whose initial diagnosis is Stage 3 (AIDS).

LATE DIAGNOSES are the subset of HIV diagnoses who are found to be Stage 3 (AIDS) within 12 months of HIV diagnosis date. Such instances are termed late diagnoses because of the advanced stage at diagnosis indicates that the person was most likely infected for many years before diagnosis.

STAGE 3 (AIDS) DIAGNOSES include two groups: 1) people newly diagnosed with HIV in 2024, found to meet the definition for Stage 3 (AIDS) at diagnosis, and with a Georgia address at or near the time of Stage 3 diagnosis; and 2) people found to meet the Stage 3 definition in 2024 with a Georgia address at or near the time of Stage 3 diagnosis, regardless of the date of their initial HIV diagnosis.

PEOPLE WITH HIV are people diagnosed by December 31, 2024, who are not known to be deceased, and for whom the most recent address available is in Georgia.

PEOPLE WITH HIV, EVER STAGE 3 are people diagnosed by December 31, 2024, who are not known to be deceased, for whom the most recent address available is in Georgia, and who met the case definition for Stage 3 (AIDS) at some point.

TRANSMISSION CATEGORIES presented in this report follow the standards created by the CDC and have been used for many years. HIV transmission categories are associated with a case using a classification system that is based on the likelihood that an HIV infection may be attributed to one or more types of exposures. The summary classification results from selecting the one most likely factor to be attributed to the HIV infection from the presumed hierarchical order of probability. For surveillance purposes, HIV and AIDS cases are counted only once in the hierarchy of transmission categories. People with more than one reported factor that can likely be attributed to the HIV infection are classified in the transmission category according to the exposure that is most likely to have resulted in transmission. The exception is men who report sexual contact with other men and injection drug use; this group makes up a separate transmission category. Individuals indicated as having “No risk identified or reported (NIR/NRR)” did not have any likely modes of exposure to HIV indicated in their case investigation.

CURRENT RESIDENCE Current address is determined using the date of the most recently entered residential address as of 12/31/2024 into eHARS. The number of people with HIV is based on current address.

VITAL STATUS: People are assumed to be alive unless otherwise documented or reported. Georgia DPH performs an annual match of the eHARS database with the Georgia Vital Records death data, the National Death Index, and the Social Security Death Index to ascertain vital status.

RATES: Denominators for population rates are based on the 2024 estimates of the Georgia resident population retrieved from the Georgia Department of Public Health, Office of Health Indicators for Planning (OHIP). Rates are presented as number of cases per 100,000 individuals. Rates are censored if associated counts are <12 because these produce unstable estimates.

PERCENTAGES: Total percentages may not add up to 100% due to rounding.

COUNTS: Counts in the gender, race and ethnicity, and age categories may not add up to the overall totals due to a small number of cases with missing information (<2 percent). In the below tables, the letter “N” is used to represent counts.

REFERENCES

1. Trickey, Adam et al. “Life expectancy after 2015 of adults with HIV on long-term antiretroviral therapy in Europe and North America: a collaborative analysis of cohort studies.” *The lancet. HIV* vol. 10,5 (2024): e295-e307. doi:10.1016/S2352-3018(23)00028-0
2. Centers for Disease Control and Prevention. *Diagnoses, Deaths, and Prevalence of HIV in the United States and 6 Territories and Freely Associated States, 2023*. <https://www.cdc.gov/hiv-data/nhss/hiv-diagnoses-deaths-and-prevalence-2025.html> Published April 2025.
3. White House Office of National AIDS Policy. *National HIV/AIDS Strategy (2022-2025)*. <https://hivgov-prod-v3.s3.amazonaws.com/s3fs-public/NHAS-2022-2025.pdf>. Published 2021.

4. Department of Public Health, Rules and regulations 290-48.11 (2003); Official Code of Georgia Annotated (O.C.G.A.) § 31-22-9.2 (2011)

5. Harrison KM, Kajese T, Hall HI, Song R. Risk factor redistribution of the national HIV/AIDS surveillance data: an alternative approach. *Public Health Rep* 2008;123:618–27.

HIV DIAGNOSES AND PEOPLE WITH HIV, GEORGIA

TABLE 1. HIV DIAGNOSES, LATE DIAGNOSES, AND STAGE 3 (AIDS) DIAGNOSES, GA, 2024

	New Diagnoses (all stages)		Late HIV Diagnoses ¹			Stage 3 (AIDS) Diagnoses ²	
	N	%*	N	%*	Row %	N	%*
Total	2,485	100.0	531	100.0	21.4	1,065	100.0
Birth Sex							
Males	1,924	77.4	413	77.8	21.5	831	78.0
Females	560	22.5	118	22.2	21.1	234	22.0
Gender Identity							
Cisgender Men	1,866	75.1	398	75	21.3	807	75.8
Cisgender Women	559	22.5	118	22.2	21.1	234	22.0
Transgender Persons/Additional Gender Identity	59	2.4	15	2.8	25.4	24	2.3
Race and Ethnicity							
Black or African American, non-Hispanic	1,568	63.1	317	59.7	20.2	705	66.2
White, non-Hispanic	379	15.3	88	16.6	23.2	151	14.2
Hispanic or Latino, Any Race	420	16.9	104	19.6	24.8	155	14.6
Asian	31	1.2	4	0.8	12.9	7	0.7
American Indian or Alaska Native	5	0.2	1	0.2	20.0	1	0.1
Native Hawaiian or Pacific Islander	0	0.0	0	0.0	0.0	0	0.0
Multiple Races	71	2.9	16	3.0	22.5	44	4.1
Age Group at Diagnosis (Years)							
<13	7	0.3	1	0.2	14.3	1	0.1
13-17	23	0.9	1	0.2	4.3	2	0.2
18-24	503	20.2	62	11.7	12.3	79	7.4
25-34	937	37.7	174	32.8	18.6	337	31.6
35-44	522	21.0	131	24.7	25.1	269	25.3
45-54	269	10.8	87	16.4	32.3	179	16.8
55-64	153	6.2	56	10.5	36.6	140	13.1
65+	71	2.9	19	3.6	26.8	58	5.4
HIV Transmission Category (Unadjusted)							
Male-to-male sexual contact (MMSC)	1,297	52.2	246	46.3	19.0	526	49.4
Heterosexual contact ³	762	30.7	218	41.1	28.6	317	29.8
Injection drug use (IDU)	42	1.7	10	1.9	23.8	33	3.1
MMSC and IDU	30	1.2	9	1.7	30.0	27	2.5
Adult Other	3	0.1	0	0.0	0.0	0	0.0
Perinatal/other pediatric ⁴	7	0.3	1	0.2	14.3	9	0.8
No risk identified or reported (NIR/NRR)	344	13.8	47	8.9	13.7	153	14.4
HIV Transmission Category (Adjusted)⁵							
Male-to-male sexual contact (MMSC)	1,501	60.4	274	51.7	18.3	608	57.1
Heterosexual contact ³	871	35.0	233	43.9	26.8	376	35.3
Injection drug use (IDU)	56	2.3	12	2.3	22.1	42	3.9
MMSC and IDU	37	1.5	10	1.9	27.0	31	2.9
Adult Other	3	0.1	0	0.0	0.0	0	0.0
Perinatal/other pediatric ⁴	7	0.3	1	0.2	14.3	9	0.8
No risk identified or reported (NIR/NRR)	10	0.4	0	0.0	0.0	0	0.0

*Percentages may not add up to 100% due to rounding and missing data. 1. Late diagnosis: stage 3 (AIDS) diagnosis within 12 months of HIV diagnosis. Row % is percent of HIV diagnoses that were late diagnoses 2. Stage 3 (AIDS): includes persons diagnosed with stage 3 in 2024, regardless of year of HIV diagnosis. 3. Heterosexual contact: Defined as sexual contact with someone of the opposite sex with known risk such as injection drug use, bisexual male (applies to females only), person with hemophilia/coagulation disorder, transfusion recipient with HIV documentation, and or person with AIDS or documented HIV. 4. Perinatal/other pediatric: cases <13 years born to a mother who is HIV-positive, and cases <13 with other risk (e.g., sexual transmission). 5. Adjusted for missing risk using multiple imputation methods. Adjusted subtotals may be different from unadjusted subtotals due to rounding.

TABLE 2. HIV DIAGNOSES, LATE DIAGNOSES, AND STAGE 3 (AIDS) DIAGNOSES AMONG CISGENDER MEN, GA, 2024

	New Diagnoses (all stages)		Late HIV Diagnoses ¹			Stage 3 (AIDS) Diagnoses ²	
	N	%*	N	%*	Row %	N	%*
Total	1,866	100.0	398	100.0	21.3	807	100.0
Race and Ethnicity							
Black or African American, non-Hispanic	1,148	61.5	232	58.3	20.2	526	65.2
White, non-Hispanic	276	14.8	65	16.3	23.6	116	14.4
Hispanic or Latino, Any Race	357	19.1	84	21.1	23.5	121	15.0
Asian	21	1.1	3	0.8	14.3	6	0.7
American Indian or Alaska Native	4	0.2	1	0.3	25.0	1	0.1
Native Hawaiian or Pacific Islander	0	0.0	0	0.0	0.0	0	0.0
Multiple Races	53	2.8	13	3.3	24.5	36	4.5
Age Group at Diagnosis (Years)							
<13	2	0.1	0	0.0	0.0	0	0.0
13-17	16	0.9	1	0.3	6.3	2	0.2
18-24	415	22.2	49	12.3	11.8	61	7.6
25-34	757	40.6	144	36.2	19.0	287	35.6
35-44	379	20.3	101	25.4	26.6	203	25.2
45-54	168	9.0	56	14.1	33.3	124	15.4
55-64	92	4.9	38	9.5	41.3	95	11.8
65+	37	2.0	9	2.3	24.3	35	4.3
HIV Transmission Category (Unadjusted)							
Male-to-male sexual contact (MMSC)	1,240	66.5	232	58.3	18.7	504	62.5
Heterosexual contact ³	308	16.5	111	27.9	36.0	145	18.0
Injection drug use (IDU)	26	1.4	6	1.5	23.1	16	2.0
MMSC and IDU	29	1.6	8	2.0	27.6	26	3.2
Adult Other	1	0.1	0	0.0	0.0	0	0.0
Perinatal/other pediatric ⁴	2	0.1	0	0.0	0.0	5	0.6
No risk identified or reported (NIR/NRR)	260	13.9	41	10.3	15.8	111	13.8
HIV Transmission Category (Adjusted)⁵							
Male-to-male sexual contact (MMSC)	1,444	77.4	260	65.4	18.0	585	72.5
Heterosexual contact ³	342	18.3	121	30.4	35.3	166	20.6
Injection drug use (IDU)	34	1.8	8	2.0	23.2	21	2.6
MMSC and IDU	36	1.9	9	2.3	24.9	30	3.7
Adult Other	1	0.1	0	0.0	0.0	0	0.0
Perinatal/other pediatric ⁴	2	0.1	0	0.0	0.0	5	0.6
No risk identified or reported (NIR/NRR)	7	0.4	0	0.0	0.0	0	0.0

*Percentages may not add up to 100% due to rounding and missing data. 1. Late diagnosis: stage 3 (AIDS) diagnosis within 12 months of HIV diagnosis. Row % is percent of HIV diagnoses that were late diagnoses 2. Stage 3 (AIDS): includes persons diagnosed with stage 3 in 2024, regardless of year of HIV diagnosis. 3. Heterosexual contact: Defined as sexual contact with someone of the opposite sex with known risk such as injection drug use, bisexual male (applies to females only), person with hemophilia/coagulation disorder, transfusion recipient with HIV documentation, and or person with AIDS or documented HIV. 4. Perinatal/other pediatric: cases <13 years born to a mother who is HIV-positive, and cases <13 with other risk (e.g., sexual transmission). 5. Adjusted for missing risk using multiple imputation methods. Adjusted subtotals may be different from unadjusted subtotals due to rounding.

TABLE 3. HIV DIAGNOSES, LATE DIAGNOSES, AND STAGE 3 (AIDS) DIAGNOSES AMONG CISGENDER WOMEN, GA, 2024

	New Diagnoses (all stages)		Late HIV Diagnoses ¹			Stage 3 (AIDS) Diagnoses ²	
	N	%*	N	%*	Row %	N	%*
Total	559	100.0	118	100.0	21.1	234	100.0
Race and Ethnicity							
Black or African American, non-Hispanic	380	68.0	75	63.6	19.7	163	69.7
White, non-Hispanic	99	17.7	22	18.6	22.2	34	14.5
Hispanic or Latino, Any Race	51	9.1	16	13.6	31.4	27	11.5
Asian	10	1.8	1	0.8	10.0	1	0.4
American Indian or Alaska Native	0	0.0	0	0.0	0.0	0	0.0
Native Hawaiian or Pacific Islander	0	0.0	0	0.0	0.0	0	0.0
Multiple Races	15	2.7	3	2.5	20.0	8	3.4
Age Group at Diagnosis (Years)							
<13	5	0.9	1	0.8	20.0	1	0.4
13-17	7	1.3	0	0.0	0.0	0	0.0
18-24	64	11.4	6	5.1	9.4	10	4.3
25-34	152	27.2	22	18.6	14.5	36	15.4
35-44	138	24.7	30	25.4	21.7	64	27.4
45-54	99	17.7	31	26.3	31.3	55	23.5
55-64	60	10.7	18	15.3	30.0	45	19.2
65+	34	6.1	10	8.5	29.4	23	9.8
HIV Transmission Category (Unadjusted)							
Heterosexual contact ³	453	81.0	107	90.7	23.6	172	73.5
Injection drug use (IDU)	16	2.9	4	3.4	25.0	17	7.3
Adult Other	2	0.4	0	0.0	0.0	0	0.0
Perinatal/other pediatric ⁴	5	0.9	1	0.8	20.0	4	1.7
No risk identified or reported (NIR/NRR)	83	14.8	6	5.1	7.2	41	17.5
HIV Transmission Category (Adjusted)⁵							
Heterosexual contact ³	528	94.4	113	95.3	21.3	209	89.4
Injection drug use (IDU)	22	4.0	5	3.8	20.4	21	8.9
Adult Other	2	0.4	0	0.0	0.0	0	0.0
Perinatal/other pediatric ⁴	5	0.9	1	0.8	20.0	4	1.7
No risk identified or reported (NIR/NRR)	2	0.4	0	0.0	0.0	0	0.0

*Percentages may not add up to 100% due to rounding and missing data. 1. Late diagnosis: stage 3 (AIDS) diagnosis within 12 months of HIV diagnosis. Row % is percent of HIV diagnoses that were late diagnoses 2. Stage 3 (AIDS): includes persons diagnosed with stage 3 in 2024, regardless of year of HIV diagnosis. 3. Heterosexual contact: Defined as sexual contact with someone of the opposite sex with known risk such as injection drug use, bisexual male (applies to females only), person with hemophilia/coagulation disorder, transfusion recipient with HIV documentation, and or person with AIDS or documented HIV. 4. Perinatal/other pediatric: cases <13 years born to a mother who is HIV-positive, and cases <13 with other risk (e.g., sexual transmission). 5. Adjusted for missing risk using multiple imputation methods. Adjusted subtotals may be different from unadjusted subtotals due to rounding.

TABLE 4. HIV DIAGNOSES, LATE DIAGNOSES, AND STAGE 3 (AIDS) DIAGNOSES AMONG TRANSGENDER PERSONS AND PERSONS WITH AN ADDITIONAL GENDER IDENTITY, GA, 2024

	New Diagnoses (all stages)		Late HIV Diagnoses ¹			Stage 3 (AIDS) Diagnoses ²	
	N	%*	N	%*	Row %	N	%*
Total	59	100.0	15	100.0	25.4	24	100.0
Birth Sex							
Males	58	98.3	15	100.0	25.9	24	100.0
Females	1	1.7	0	0.0	0.0	0	0.0
Gender Identity							
Transgender Women	54	91.5	14	93.3	25.9	23	95.8
Transgender Men	1	1.7	0	0.0	0.0	0	0.0
Additional Gender Identity	4	6.8	1	6.7	25.0	1	4.2
Race and Ethnicity							
Black or African American, non-Hispanic	40	67.8	10	66.7	25.0	16	66.7
White, non-Hispanic	4	6.8	1	6.7	25.0	1	4.2
Hispanic or Latino, Any Race	11	18.6	4	26.7	36.4	7	29.2
Asian	0	0.0	0	0.0	0.0	0	0.0
American Indian or Alaska Native	1	1.7	0	0.0	0.0	0	0.0
Native Hawaiian or Pacific Islander	0	0.0	0	0.0	0.0	0	0.0
Multiple Races	3	5.1	0	0.0	0.0	0	0.0
Age Group at Diagnosis (Years)							
<13	0	0.0	0	0.0	0.0	0	0.0
13-17	0	0.0	0	0.0	0.0	0	0.0
18-24	24	40.7	7	46.7	29.2	8	33.3
25-34	28	47.5	8	53.3	28.6	14	58.3
35-44	4	6.8	0	0.0	0.0	2	8.3
45-54	2	3.4	0	0.0	0.0	0	0.0
55-64	1	1.7	0	0.0	0.0	0	0.0
65+	0	0.0	0	0.0	0.0	0	0.0
HIV Transmission Category (Unadjusted)							
Sexual contact ³	58	98.3	14	93.3	24.6	22	91.7
Injection drug use (IDU)	0	0.0	0	0.0	0.0	0	0.0
Sexual contact and IDU	1	1.7	1	6.7	100.0	1	4.2
Adult Other	0	0.0	0	0.0	0.0	0	0.0
Perinatal/other pediatric	0	0.0	0	0.0	0.0	0	0.0
No risk identified or reported (NIR/NRR)	0	0.0	0	0.0	0.0	1	4.2
HIV Transmission Category (Adjusted)⁴							
Sexual contact ³	58	98.3	14	93.3	24.6	23	95.0
Injection drug use (IDU)	0	0.0	0	0.0	0.0	0	0.0
Sexual contact and IDU	1	1.7	1	6.7	100.0	1	4.6
Adult Other	0	0.0	0	0.0	0.0	0	0.0
Perinatal/other pediatric	0	0.0	0	0.0	0.0	0	0.0
No risk identified or reported (NIR/NRR)	0	0.0	0	0.0	0.0	0	0.0

* Percentages may not add up to 100% due to rounding. 1. Late diagnosis: stage 3 (AIDS) diagnosis within 12 months of HIV diagnosis. 2. Stage 3 (AIDS): includes persons diagnosed with stage 3 in 2024, regardless of year of HIV diagnosis. 3. Includes sex with male and probable heterosexual transmission as noted in medical chart. 4. Adjusted for missing risk using multiple imputation methods. Adjusted subtotals may be different from unadjusted subtotals due to rounding.

TABLE 5. HIV DIAGNOSES, LATE DIAGNOSES, AND STAGE 3 (AIDS) DIAGNOSES, AMONG BLACK OR AFRICAN AMERICAN, NON-HISPANIC, INDIVIDUALS, GA, 2024

	New Diagnoses (all stages)		Late HIV Diagnoses ¹			Stage 3 (AIDS) Diagnoses ²	
	N	%*	N	%*	Row %	N	%*
Total	1,568	100.0	317	100.0	20.2	705	100.0
Birth Sex							
Males	1,188	75.8	242	76.3	20.4	542	76.9
Females	380	24.2	75	23.7	19.7	163	23.1
Gender Identity							
Cisgender Men	1,148	73.2	232	73.2	20.2	526	74.6
Cisgender Women	380	24.2	75	23.7	19.7	163	23.1
Transgender Persons/Additional Gender Identity	40	2.6	10	3.2	25.0	16	2.3
Age Group at Diagnosis (Years)							
<13	7	0.4	1	0.3	14.3	1	0.1
13-17	20	1.3	1	0.3	5.0	2	0.3
18-24	362	23.1	42	13.2	11.6	55	7.8
25-34	568	36.2	95	30.0	16.7	221	31.3
35-44	321	20.5	82	25.9	25.5	182	25.8
45-54	153	9.8	53	16.7	34.6	117	16.6
55-64	92	5.9	31	9.8	33.7	87	12.3
65+	45	2.9	12	3.8	26.7	40	5.7
HIV Transmission Category (Unadjusted)							
Male-to-male sexual contact (MMSC)	791	50.4	148	46.7	18.7	346	49.1
Heterosexual contact ³	521	33.2	138	43.5	26.5	220	31.2
Injection drug use (IDU)	16	1.0	3	0.9	18.8	17	2.4
MMSC and IDU	6	0.4	1	0.3	16.7	10	1.4
Adult Other	2	0.1	0	0.0	0.0	0	0.0
Perinatal/other pediatric ⁴	7	0.4	1	0.3	14.3	7	1.0
No risk identified or reported (NIR/NRR)	225	14.3	26	8.2	11.6	105	14.9
HIV Transmission Category (Adjusted)⁵							
Male-to-male sexual contact (MMSC)	923	58.9	161	50.9	17.5	398	56.5
Heterosexual contact ³	600	38.2	148	46.8	24.7	264	37.4
Injection drug use (IDU)	26	1.7	5	1.5	17.6	24	3.4
MMSC and IDU	10	0.7	2	0.5	16.7	12	1.8
Adult Other	2	0.1	0	0.0	0.0	0	0.0
Perinatal/other pediatric ⁴	7	0.4	1	0.3	14.3	7	1.0
No risk identified or reported (NIR/NRR)	0	0.0	0	0.0	0.0	0	0.0

*Percentages may not add up to 100% due to rounding and missing data. 1. Late diagnosis: stage 3 (AIDS) diagnosis within 12 months of HIV diagnosis. Row % is percent of HIV diagnoses that were late diagnoses 2. Stage 3 (AIDS): includes persons diagnosed with stage 3 in 2024, regardless of year of HIV diagnosis. 3. Heterosexual contact: Defined as sexual contact with someone of the opposite sex with known risk such as injection drug use, bisexual male (applies to females only), person with hemophilia/coagulation disorder, transfusion recipient with HIV documentation, and or person with AIDS or documented HIV. 4. Perinatal/other pediatric: cases <13 years born to a mother who is HIV-positive, and cases <13 with other risk (e.g., sexual transmission). 5. Adjusted for missing risk using multiple imputation methods. Adjusted subtotals may be different from unadjusted subtotals due to rounding.

TABLE 6. HIV DIAGNOSES, LATE DIAGNOSES, AND STAGE 3 (AIDS) DIAGNOSES, AMONG WHITE, NON-HISPANIC, INDIVIDUALS, GA, 2024

	New Diagnoses (all stages)		Late HIV Diagnoses ¹			Stage 3 (AIDS) Diagnoses ²	
	N	%*	N	%*	Row %	N	%*
Total	379	100.0	88	100.0	23.2	151	100.0
Birth Sex							
Males	279	73.6	66	75.0	23.7	117	77.5
Females	100	26.4	22	25.0	22.0	34	22.5
Gender Identity							
Cisgender Men	276	72.8	65	73.9	23.6	116	76.8
Cisgender Women	99	26.1	22	25.0	22.2	34	22.5
Transgender Persons/Additional Gender Identity	4	1.1	1	1.1	25.0	1	0.7
Age Group at Diagnosis (Years)							
<13	0	0.0	0	0.0	0.0	0	0.0
13-17	1	0.3	0	0.0	0.0	0	0.0
18-24	39	10.3	6	6.8	15.4	5	3.3
25-34	135	35.6	24	27.3	17.8	38	25.2
35-44	79	20.8	19	21.6	24.1	31	20.5
45-54	75	19.8	24	27.3	32.0	38	25.2
55-64	35	9.2	13	14.8	37.1	31	20.5
65+	15	4.0	2	2.3	13.3	8	5.3
HIV Transmission Category (Unadjusted)							
Male-to-male sexual contact (MMSC)	169	44.6	32	36.4	18.9	65	43.0
Heterosexual contact ³	114	30.1	35	39.8	30.7	43	28.5
Injection drug use (IDU)	20	5.3	6	6.8	30.0	11	7.3
MMSC and IDU	15	4.0	6	6.8	40.0	13	8.6
Adult Other	0	0.0	0	0.0	0.0	0	0.0
Perinatal/other pediatric ⁴	0	0.0	0	0.0	0.0	0	0.0
No risk identified or reported (NIR/NRR)	61	16.1	9	10.2	14.8	19	12.6
HIV Transmission Category (Adjusted)⁵							
Male-to-male sexual contact (MMSC)	206	54.4	40	45.5	19.4	78	51.5
Heterosexual contact ³	134	35.4	36	40.8	26.8	48	31.7
Injection drug use (IDU)	22	5.9	6	6.8	26.9	11	7.5
MMSC and IDU	17	4.4	6	6.9	37.0	14	9.2
Adult Other	0	0.0	0	0.0	0.0	0	0.0
Perinatal/other pediatric ⁴	0	0.0	0	0.0	0.0	0	0.0
No risk identified or reported (NIR/NRR)	0	0.0	0	0.0	0.0	0	0.0

*Percentages may not add up to 100% due to rounding and missing data. 1. Late diagnosis: stage 3 (AIDS) diagnosis within 12 months of HIV diagnosis. Row % is percent of HIV diagnoses that were late diagnoses 2. Stage 3 (AIDS): includes persons diagnosed with stage 3 in 2024, regardless of year of HIV diagnosis. 3. Heterosexual contact: Defined as sexual contact with someone of the opposite sex with known risk such as injection drug use, bisexual male (applies to females only), person with hemophilia/coagulation disorder, transfusion recipient with HIV documentation, and or person with AIDS or documented HIV. 4. Perinatal/other pediatric: cases <13 years born to a mother who is HIV-positive, and cases <13 with other risk (e.g., sexual transmission). 5. Adjusted for missing risk using multiple imputation methods. Adjusted subtotals may be different from unadjusted subtotals due to rounding.

TABLE 7. HIV DIAGNOSES, LATE DIAGNOSES, AND STAGE 3 (AIDS) DIAGNOSES, AMONG HISPANIC OR LATINO (OF ANY RACE) INDIVIDUALS, GA, 2024

	New Diagnoses (all stages)		Late HIV Diagnoses ¹			Stage 3 (AIDS) Diagnoses ²	
	N	%*	N	%*	Row %	N	%*
Total	420	100.0	104	100.0	24.8	155	100.0
Birth Sex							
Males	368	87.6	88	84.6	23.9	128	82.6
Females	51	12.1	16	15.4	31.4	27	17.4
Gender Identity							
Cisgender Men	357	85.0	84	80.8	23.5	121	78.1
Cisgender Women	51	12.1	16	15.4	31.4	27	17.4
Transgender Persons/Additional Gender Identity	11	2.6	4	3.8	36.4	7	4.5
Age Group at Diagnosis (Years)							
<13	0	0.0	0	0.0	0.0	0	0.0
13-17	2	0.5	0	0.0	0.0	0	0.0
18-24	81	19.3	13	12.5	16.0	18	11.6
25-34	192	45.7	45	43.3	23.4	62	40.0
35-44	91	21.7	23	22.1	25.3	39	25.2
45-54	29	6.9	8	7.7	27.6	16	10.3
55-64	20	4.8	11	10.6	55.0	14	9.0
65+	5	1.2	4	3.8	80.0	6	3.9
HIV Transmission Category (Unadjusted)							
Male-to-male sexual contact (MMSC)	276	65.7	56	53.8	20.3	84	54.2
Heterosexual contact ³	94	22.4	36	34.6	38.3	43	27.7
Injection drug use (IDU)	3	0.7	0	0.0	0.0	3	1.9
MMSC and IDU	5	1.2	2	1.9	40.0	4	2.6
Adult Other	1	0.2	0	0.0	0.0	0	0.0
Perinatal/other pediatric ⁴	0	0.0	0	0.0	0.0	1	0.6
No risk identified or reported (NIR/NRR)	41	9.8	10	9.6	24.4	20	12.9
HIV Transmission Category (Adjusted)⁵							
Male-to-male sexual contact (MMSC)	306	72.8	62	59.4	20.2	98	63.1
Heterosexual contact ³	102	24.2	40	38.0	38.9	48	30.9
Injection drug use (IDU)	4	1.0	1	0.5	11.9	4	2.5
MMSC and IDU	6	1.5	2	2.1	34.9	5	2.9
Adult Other	1	0.2	0	0.0	0.0	0	0.0
Perinatal/other pediatric ⁴	0	0.0	0	0.0	0.0	1	0.6
No risk identified or reported (NIR/NRR)	1	0.2	0	0.0	0.0	0	0.0

*Percentages may not add up to 100% due to rounding and missing data. 1. Late diagnosis: stage 3 (AIDS) diagnosis within 12 months of HIV diagnosis. Row % is percent of HIV diagnoses that were late diagnoses 2. Stage 3 (AIDS): includes persons diagnosed with stage 3 in 2024, regardless of year of HIV diagnosis. 3. Heterosexual contact: Defined as sexual contact with someone of the opposite sex with known risk such as injection drug use, bisexual male (applies to females only), person with hemophilia/coagulation disorder, transfusion recipient with HIV documentation, and or person with AIDS or documented HIV. 4. Perinatal/other pediatric: cases <13 years born to a mother who is HIV-positive, and cases <13 with other risk (e.g., sexual transmission). 5. Adjusted for missing risk using multiple imputation methods. Adjusted subtotals may be different from unadjusted subtotals due to rounding.

TABLE 8. PEOPLE WITH HIV (PWH), AND PWH EVER STAGE 3 (AIDS), GA, 2024

	PWH		Ever Stage 3 (AIDS) ¹	
	N	%*	N	%*
Total	66,898	100.0	31,602	100.0
Birth Sex				
Males	51,574	77.1	24,037	76.1
Females	15,277	22.8	7,553	23.9
Gender Identity				
Cisgender Men	50,448	75.4	23,574	74.6
Cisgender Women	15,231	22.8	7,532	23.8
Transgender Persons/Additional Gender Identity	1,172	1.8	484	1.5
Race and Ethnicity				
Black or African American, non-Hispanic	44,592	66.7	21,517	68.1
White, non-Hispanic	10,530	15.7	4,862	15.4
Hispanic or Latino, Any Race	6,549	9.8	2,839	9.0
Asian	383	0.6	133	0.4
American Indian or Alaska Native	23	0.0	9	0.0
Native Hawaiian or Pacific Islander	10	0.0	5	0.0
Multiple Races	4,056	6.1	2,125	6.7
Current Age Group (Years)				
<13	76	0.1	24	0.1
13-17	86	0.1	17	0.1
18-24	2,020	3.0	299	0.9
25-34	12,371	18.5	3,186	10.1
35-44	16,173	24.2	6,605	20.9
45-54	13,295	19.9	7,149	22.6
55-64	14,822	22.2	9,165	29.0
65+	8,053	12.0	5,156	16.3
HIV Transmission Category (Unadjusted)				
Male-to-male sexual contact (MMSC)	35,719	53.4	15,926	50.4
Heterosexual contact ²	12,034	18.0	6,284	19.9
Injection drug use (IDU)	2,299	3.4	1,571	5.0
MMSC and IDU	2,122	3.2	1,278	4.0
Adult Other ³	668	1.0	340	1.1
Perinatal/other pediatric ⁴	69	0.1	23	0.1
No risk identified or reported (NIR/NRR)	13,987	20.9	6,180	19.6
HIV Transmission Category (Adjusted)⁵				
Male-to-male sexual contact (MMSC)	42,124	63.0	18,684	59.1
Heterosexual contact ²	17,481	26.1	8,827	27.9
Injection drug use (IDU)	3,241	4.8	2,103	6.7
MMSC and IDU	2,564	3.8	1,513	4.8
Adult Other ³	668	1.0	340	1.1
Perinatal/other pediatric ⁴	69	0.1	23	0.1
No risk identified or reported (NIR/NRR)	751	1.1	112	0.4

* Percentages may not add up to 100% due to rounding. 1. Persons whose infection was ever classified as stage 3 (AIDS). 2. Heterosexual contact: Defined as sexual contact with someone of the opposite sex with known risk such as injection drug use, bisexual male (applies to females only), person with hemophilia/coagulation disorder, transfusion recipient with HIV documentation, and or person with AIDS or documented HIV. 3. Includes perinatal exposure current age ≥ 13, hemophilia, blood transfusion. 4. Perinatal/other pediatric: cases <13 born to a mother who is HIV-positive and cases <13 with other risk (e.g., sexual transmission). 5. Adjusted for missing risk using multiple imputation methods.

TABLE 9. PEOPLE WITH HIV (PWH) AND PWH EVER STAGE 3 (AIDS), CISGENDER MEN, GA, 2024

	PWH		Ever Stage 3 (AIDS) ¹	
	N	%*	N	%*
Total	50,448	100.0	23,574	100.0
Race and Ethnicity				
Black or African American, non-Hispanic	32,272	64.0	15,324	65.0
White, non-Hispanic	8,895	17.6	4,165	17.7
Hispanic or Latino, Any Race	5,289	10.5	2,263	9.6
Asian	288	0.6	105	0.4
American Indian or Alaska Native	19	0.0	8	0.0
Native Hawaiian or Pacific Islander	7	0.0	4	0.0
Multiple Races	3,145	6.2	1,625	6.9
Current Age Group (Years)				
<13	32	0.1	11	0.0
13-17	47	0.1	12	0.1
18-24	1,610	3.2	232	1.0
25-34	10,283	20.4	2,611	11.1
35-44	12,692	25.2	5,188	22.0
45-54	9,322	18.5	5,035	21.4
55-64	10,855	21.5	6,807	28.9
65+	5,605	11.1	3,677	15.6
HIV Transmission Category (Unadjusted)				
Male-to-male sexual contact (MMSC)	34,669	68.7	15,502	65.8
Heterosexual contact ²	3,323	6.6	1,895	8.0
Injection drug use (IDU)	1,237	2.5	858	3.6
MMSC and IDU	2,072	4.1	1,251	5.3
Adult Other ³	313	0.6	154	0.7
Perinatal/other pediatric ⁴	29	0.1	10	0.0
No risk identified or reported (NIR/NRR)	8,805	17.5	3,904	16.6
HIV Transmission Category (Adjusted)⁵				
Male-to-male sexual contact (MMSC)	41,062	81.4	18,256	77.4
Heterosexual contact ²	4,391	8.7	2,503	10.6
Injection drug use (IDU)	1,625	3.2	1,088	4.6
MMSC and IDU	2,513	5.0	1,486	6.3
Adult Other ³	313	0.6	154	0.7
Perinatal/other pediatric ⁴	29	0.1	10	0.0
No risk identified or reported (NIR/NRR)	515	1.0	77	0.3

* Percentages may not add up to 100% due to rounding. 1. Persons whose infection was ever classified as stage 3 (AIDS). 2. Heterosexual contact: Defined as sexual contact with someone of the opposite sex with known risk such as injection drug use, bisexual male (applies to females only), person with hemophilia/coagulation disorder, transfusion recipient with HIV documentation, and or person with AIDS or documented HIV. 3. Includes perinatal exposure current age \geq 13, hemophilia, blood transfusion. 4. Perinatal/other pediatric: cases <13 born to a mother who is HIV-positive and cases <13 with other risk (e.g., sexual transmission). 5. Adjusted for missing risk using multiple imputation methods.

TABLE 10. PEOPLE WITH HIV (PWH) AND PWH EVER STAGE 3 (AIDS), CISGENDER WOMEN, GA, 2024

	PWH		Ever Stage 3 (AIDS) ¹	
	N	%*	N	%*
Total	15,231	100.0	7,532	100.0
Race and Ethnicity				
Black or African American, non-Hispanic	11,442	75.1	5,825	77.3
White, non-Hispanic	1,570	10.3	678	9.0
Hispanic or Latino, Any Race	1,098	7.2	507	6.7
Asian	93	0.6	26	0.3
American Indian or Alaska Native	3	0.0	1	0.0
Native Hawaiian or Pacific Islander	3	0.0	1	0.0
Multiple Races	830	5.4	470	6.2
Current Age Group (Years)				
<13	44	0.3	13	0.2
13-17	38	0.2	5	0.1
18-24	337	2.2	51	0.7
25-34	1,674	11.0	457	6.1
35-44	3,052	20.0	1,228	16.3
45-54	3,792	24.9	2,014	26.7
55-64	3,874	25.4	2,303	30.6
65+	2,420	15.9	1,461	19.4
HIV Transmission Category (Unadjusted)				
Heterosexual contact ²	8,687	57.0	4,377	58.1
Injection drug use (IDU)	1,055	6.9	709	9.4
Adult Other ³	348	2.3	182	2.4
Perinatal/other pediatric ⁴	40	0.3	13	0.2
No risk identified or reported (NIR/NRR)	5,101	33.5	2,251	29.9
HIV Transmission Category (Adjusted)⁵				
Heterosexual contact ²	13,048	85.7	6,305	83.7
Injection drug use (IDU)	1,606	10.5	1,009	13.4
Adult Other ³	348	2.3	182	2.4
Perinatal/other pediatric ⁴	40	0.3	13	0.2
No risk identified or reported (NIR/NRR)	189	1.2	23	0.3

* Percentages may not add up to 100% due to rounding. 1. Persons whose infection was ever classified as stage 3 (AIDS). 2. Heterosexual contact: Defined as sexual contact with someone of the opposite sex with known risk such as injection drug use, bisexual male (applies to females only), person with hemophilia/coagulation disorder, transfusion recipient with HIV documentation, and or person with AIDS or documented HIV. 3. Includes perinatal exposure current age ≥ 13, hemophilia, blood transfusion. 4. Perinatal/other pediatric: cases <13 born to a mother who is HIV-positive and cases <13 with other risk (e.g., sexual transmission). 5. Adjusted for missing risk using multiple imputation methods.

TABLE 11. PEOPLE WITH HIV (PWH), AND PWH EVER STAGE 3 (AIDS), TRANSGENDER PERSONS OR PERSONS WITH AN ADDITIONAL GENDER IDENTITY, GA, 2024

	PWH		Ever Stage 3 (AIDS) ¹	
	N	%*	N	%*
Total	1,172	100.0	484	100.0
Birth Sex				
Males	1,126	96.1	463	95.7
Females	46	3.9	21	4.3
Gender Identity				
Transgender Women	1,081	92.2	443	91.5
Transgender Men	37	3.2	13	2.7
Additional Gender Identity	54	4.6	28	5.8
Race and Ethnicity				
Black or African American, non-Hispanic	867	74.0	364	75.2
White, non-Hispanic	62	5.3	19	3.9
Hispanic or Latino, Any Race	159	13.6	69	14.3
Asian	2	0.2	2	0.4
American Indian or Alaska Native	1	0.1	0	0.0
Native Hawaiian or Pacific Islander	0	0.0	0	0.0
Multiple Races	81	6.9	30	6.2
Current Age Group (Years)				
<13	0	0.0	0	0.0
13-17	1	0.1	0	0.0
18-24	72	6.1	16	3.3
25-34	412	35.2	118	24.4
35-44	423	36.1	187	38.6
45-54	158	13.5	94	19.4
55-64	82	7.0	53	11.0
65+	24	2.0	16	3.3
HIV Transmission Category (Unadjusted)				
Sexual contact ²	1,074	92.0	436	90.0
Injection drug use (IDU)	7	0.6	4	0.8
Sexual contact and IDU	50	4.3	27	5.6
Adult Other	7	0.6	4	0.8
Perinatal/other pediatric	0	0.0	0	0.0
No risk identified or reported (NIR/NRR)	34	2.9	13	2.7
HIV Transmission Category (Adjusted)³				
Sexual contact ²	1,104	94.0	447	92.0
Injection drug use (IDU)	10	0.9	6	1.2
Sexual contact and IDU	51	4.3	28	5.7
Adult Other	7	0.6	4	0.8
Perinatal/other pediatric	0	0.0	0	0.0
No risk identified or reported (NIR/NRR)	0	0.0	0	0.0

* Percentages may not add up to 100% due to rounding. 1. Persons whose infection was ever classified as stage 3 (AIDS). 2. Includes sex with male and probable heterosexual transmission as noted in medical chart 3. Adjusted for missing risk using multiple imputation methods. Adjusted subtotals may be different from unadjusted subtotals due to rounding.

TABLE 12. PEOPLE WITH HIV (PWH), AND PWH EVER STAGE 3 (AIDS), AMONG BLACK OR AFRICAN AMERICAN, NON-HISPANIC, INDIVIDUALS, GA, 2024

	PWH		Ever Stage 3 (AIDS) ¹	
	N	%*	N	%*
Total	44,592	100.0	21,517	100.0
Birth Sex				
Males	33,109	74.2	15,674	72.8
Females	11,472	25.7	5,839	27.1
Gender Identity				
Cisgender Men	32,272	72.4	15,324	71.2
Cisgender Women	11,442	25.7	5,825	27.1
Transgender Persons/Additional Gender Identity	867	1.9	364	1.7
Current Age Group (Years)				
<13	57	0.1	23	0.1
13-17	63	0.1	15	0.1
18-24	1,534	3.4	225	1.0
25-34	9,047	20.3	2,462	11.4
35-44	11,373	25.5	4,874	22.7
45-54	8,524	19.1	4,800	22.3
55-64	8,998	20.2	5,797	26.9
65+	4,995	11.2	3,320	15.4
HIV Transmission Category (Unadjusted)				
Male-to-male sexual contact (MMSC)	23,044	51.7	10,239	47.6
Heterosexual contact ²	9,232	20.7	4,880	22.7
Injection drug use (IDU)	1,468	3.3	1,058	4.9
MMSC and IDU	1,069	2.4	700	3.3
Adult Other ³	472	1.1	243	1.1
Perinatal/other pediatric ⁴	55	0.1	22	0.1
No risk identified or reported (NIR/NRR)	9,252	20.7	4,375	20.3
HIV Transmission Category (Adjusted)⁵				
Male-to-male sexual contact (MMSC)	27,081	60.7	12,012	55.8
Heterosexual contact ²	13,496	30.3	6,930	32.2
Injection drug use (IDU)	2,151	4.8	1,462	6.8
MMSC and IDU	1,324	3.0	843	3.9
Adult Other ³	472	1.1	243	1.1
Perinatal/other pediatric ⁴	55	0.1	22	0.1
No risk identified or reported (NIR/NRR)	13	0.0	5	0.0

* Percentages may not add up to 100% due to rounding. 1. Persons whose infection was ever classified as stage 3 (AIDS). 2. Heterosexual contact: Defined as sexual contact with someone of the opposite sex with known risk such as injection drug use, bisexual male (applies to females only), person with hemophilia/coagulation disorder, transfusion recipient with HIV documentation, and or person with AIDS or documented HIV. 3. Includes perinatal exposure current age ≥ 13, hemophilia, blood transfusion. 4. Perinatal/other pediatric: cases <13 born to a mother who is HIV-positive and cases <13 with other risk (e.g., sexual transmission). 5. Adjusted for missing risk using multiple imputation methods.

TABLE 13. PEOPLE WITH HIV (PWH), AND PWH EVER STAGE 3 (AIDS), AMONG WHITE, NON-HISPANIC, INDIVIDUALS, GA, 2024

	PWH		Ever Stage 3 (AIDS) ¹	
	N	%*	N	%*
Total	10,530	100.0	4,862	100.0
Birth Sex				
Males	8,951	85.0	4,182	86.0
Females	1,576	15.0	680	14.0
Gender Identity				
Cisgender Men	8,895	84.5	4,165	85.7
Cisgender Women	1,570	14.9	678	13.9
Transgender Persons/Additional Gender Identity	62	0.6	19	0.4
Current Age Group (Years)				
<13	0	0.0	0	0.0
13-17	8	0.1	0	0.0
18-24	138	1.3	18	0.4
25-34	1,087	10.3	203	4.2
35-44	1,723	16.4	543	11.2
45-54	2,112	20.1	959	19.7
55-64	3,456	32.8	1,944	40.0
65+	2,006	19.1	1,195	24.6
HIV Transmission Category (Unadjusted)				
Male-to-male sexual contact (MMSC)	6,161	58.5	2,943	60.5
Heterosexual contact ²	1,023	9.7	484	10.0
Injection drug use (IDU)	471	4.5	273	5.6
MMSC and IDU	627	6.0	340	7.0
Adult Other ³	60	0.6	33	0.7
Perinatal/other pediatric ⁴	0	0.0	0	0.0
No risk identified or reported (NIR/NRR)	2,188	20.8	789	16.2
HIV Transmission Category (Adjusted)⁵				
Male-to-male sexual contact (MMSC)	7,536	71.6	3,437	70.7
Heterosexual contact ²	1,580	15.0	672	13.8
Injection drug use (IDU)	620	5.9	335	6.9
MMSC and IDU	732	7.0	385	7.9
Adult Other ³	60	0.6	33	0.7
Perinatal/other pediatric ⁴	0	0.0	0	0.0
No risk identified or reported (NIR/NRR)	3	0.0	0	0.0

* Percentages may not add up to 100% due to rounding. 1. Persons whose infection was ever classified as stage 3 (AIDS). 2. Heterosexual contact: Defined as sexual contact with someone of the opposite sex with known risk such as injection drug use, bisexual male (applies to females only), person with hemophilia/coagulation disorder, transfusion recipient with HIV documentation, and or person with AIDS or documented HIV. 3. Includes perinatal exposure current age ≥ 13, hemophilia, blood transfusion. 4. Perinatal/other pediatric: cases <13 born to a mother who is HIV-positive and cases <13 with other risk (e.g., sexual transmission). 5. Adjusted for missing risk using multiple imputation methods.

TABLE 14. PEOPLE WITH HIV (PWH), AND PWH EVER STAGE 3 (AIDS), AMONG HISPANIC OR LATINO (OF ANY RACE) INDIVIDUALS, GA, 2024

	PWH		Ever Stage 3 (AIDS) ¹	
	N	%*	N	%*
Total	6,549	100.0	2,839	100.0
Birth Sex				
Males	5,441	83.1	2,330	82.1
Females	1,105	16.9	509	17.9
Gender Identity				
Cisgender Men	5,289	80.8	2,263	79.7
Cisgender Women	1,098	16.8	507	17.9
Transgender Persons/Additional Gender Identity	159	2.4	69	2.4
Current Age Group (Years)				
<13	6	0.1	0	0.0
13-17	7	0.1	2	0.1
18-24	233	3.6	40	1.4
25-34	1,458	22.3	331	11.7
35-44	1,855	28.3	696	24.5
45-54	1,453	22.2	777	27.4
55-64	1,120	17.1	715	25.2
65+	417	6.4	278	9.8
HIV Transmission Category (Unadjusted)				
Male-to-male sexual contact (MMSC)	3,847	58.7	10,239	47.6
Heterosexual contact ²	1,057	16.1	4,880	22.7
Injection drug use (IDU)	194	3.0	1,058	4.9
MMSC and IDU	211	3.2	700	3.3
Adult Other ³	76	1.2	243	1.1
Perinatal/other pediatric ⁴	6	0.1	22	0.1
No risk identified or reported (NIR/NRR)	1,158	17.7	4,375	20.3
HIV Transmission Category (Adjusted)⁵				
Male-to-male sexual contact (MMSC)	4,525	69.1	1,789	63.0
Heterosexual contact ²	1,424	21.7	704	24.8
Injection drug use (IDU)	256	3.9	169	6.0
MMSC and IDU	259	4.0	146	5.2
Adult Other ³	76	1.2	31	1.1
Perinatal/other pediatric ⁴	6	0.1	0	0.0
No risk identified or reported (NIR/NRR)	3	0.0	0	0.0

* Percentages may not add up to 100% due to rounding. 1. Persons whose infection was ever classified as stage 3 (AIDS). 2. Heterosexual contact: Defined as sexual contact with someone of the opposite sex with known risk such as injection drug use, bisexual male (applies to females only), person with hemophilia/coagulation disorder, transfusion recipient with HIV documentation, and or person with AIDS or documented HIV. 3. Includes perinatal exposure current age \geq 13, hemophilia, blood transfusion. 4. Perinatal/other pediatric: cases <13 born to a mother who is HIV-positive and cases <13 with other risk (e.g., sexual transmission). 5. Adjusted for missing risk using multiple imputation methods.

HIV DIAGNOSES AND PEOPLE WITH HIV BY PUBLIC HEALTH DISTRICT

TABLE 15. HIV DIAGNOSES AND PEOPLE WITH HIV (PWH), DISTRICT 1-1 (ROME), GA, 2024

	HIV Diagnoses		PWH	
	N	%*	N	%*
Total	77	100.0	1,563	100.0
Birth Sex				
Males	58	75.3	1,141	73.0
Females	19	24.7	422	27.0
Gender Identity				
Cisgender Men	58	75.3	1,127	72.1
Cisgender Women	19	24.7	421	26.9
Transgender Persons/Additional Gender Identity	0	0.0	15	1.0
Race and Ethnicity				
Black or African American, non-Hispanic	28	36.4	636	40.7
White, non-Hispanic	36	46.8	646	41.3
Hispanic or Latino, Any Race	8	10.4	170	10.9
Asian	0	0.0	10	0.6
American Indian or Alaska Native	0	0.0	0	0.0
Native Hawaiian or Pacific Islander	0	0.0	0	0.0
Multiple Races	5	6.5	87	5.6
Age Group (at diagnosis for diagnosed, current for PWH)				
<13	0	0.0	<5	---
13-17	<5	---	<5	---
18-24	12	15.6	39	2.5
25-34	29	37.7	266	17.0
35-44	20	26.0	327	20.9
45-54	7	9.1	340	21.8
55-64	7	9.1	371	23.7
65+	<5	---	216	13.8
HIV Transmission Category (Unadjusted)				
Male-to-male sexual contact (MMSC)	28	36.4	703	45.0
Heterosexual contact ¹	31	40.3	305	19.5
Injection drug use (IDU)	5	6.5	88	5.6
MMSC and IDU	<5	---	83	5.3
Adult Other ²	0	0.0	23	1.5
Perinatal/other pediatric ³	0	0.0	<5	---
No risk identified or reported (NIR/NRR)	9	11.7	360	23.0
HIV Transmission Category (Adjusted)⁴				
Male-to-male sexual contact (MMSC)	35	44.8	872	55.8
Heterosexual contact ¹	33	42.6	443	28.4
Injection drug use (IDU)	5	6.9	115	7.3
MMSC and IDU	<5	---	95	6.1
Adult Other ²	0	0.0	23	1.5
Perinatal/other pediatric ³	0	0.0	<5	---
No risk identified or reported (NIR/NRR)	0	0.0	14	0.9

* Percentages may not add up to 100% due to rounding. 1. Heterosexual contact: Defined as sexual contact with someone of the opposite sex with known risk such as injection drug use, bisexual male (applies to females only), person with hemophilia/coagulation disorder, transfusion recipient with HIV documentation, and or person with AIDS or documented HIV. 2. Includes perinatal exposure current age ≥ 13 ; hemophilia, blood transfusion. 3. Perinatal/other pediatric: cases <13 born to a mother who is HIV-positive, and cases <13 with other risk (e.g., sexual transmission). 4. Adjusted for missing risk using multiple imputation methods. Adjusted subtotals may be different from unadjusted subtotals due to rounding.

TABLE 16. HIV DIAGNOSES AND PEOPLE WITH HIV (PWH), DISTRICT 1-2 (DALTON), GA, 2024

	HIV Diagnoses		PWH	
	N	%*	N	%*
Total	19	100.0	872	100.0
Birth Sex				
Males	12	63.2	691	79.2
Females	7	36.8	180	20.6
Gender Identity				
Cisgender Men	12	63.2	679	77.9
Cisgender Women	7	36.8	179	20.5
Transgender Persons/Additional Gender Identity	0	0.0	13	1.5
Race and Ethnicity				
Black or African American, non-Hispanic	<5	---	189	21.7
White, non-Hispanic	8	42.1	445	51.0
Hispanic or Latino, Any Race	6	31.6	170	19.5
Asian	0	0.0	<5	---
American Indian or Alaska Native	0	0.0	0	0.0
Native Hawaiian or Pacific Islander	0	0.0	0	0.0
Multiple Races	<5	---	53	6.1
Age Group (at diagnosis for diagnosed, current for PWH)				
<13	0	0.0	<5	---
13-17	0	0.0	<5	---
18-24	<5	---	11	1.3
25-34	10	52.6	143	16.4
35-44	<5	---	181	20.8
45-54	<5	---	188	21.6
55-64	<5	---	219	25.1
65+	<5	---	126	14.4
HIV Transmission Category (Unadjusted)				
Male-to-male sexual contact (MMSC)	10	52.6	442	50.7
Heterosexual contact ¹	6	31.6	152	17.4
Injection drug use (IDU)	<5	---	40	4.6
MMSC and IDU	0	0.0	40	4.6
Adult Other ²	0	0.0	7	0.8
Perinatal/other pediatric ³	0	0.0	<5	---
No risk identified or reported (NIR/NRR)	<5	---	190	21.8
HIV Transmission Category (Adjusted)⁴				
Male-to-male sexual contact (MMSC)	12	61.6	543	62.3
Heterosexual contact ¹	6	33.2	210	24.1
Injection drug use (IDU)	<5	---	50	5.8
MMSC and IDU	0	0.0	48	5.6
Adult Other ²	0	0.0	7	0.8
Perinatal/other pediatric ³	0	0.0	<5	---
No risk identified or reported (NIR/NRR)	0	0.0	12	1.4

* Percentages may not add up to 100% due to rounding. 1. Heterosexual contact: Defined as sexual contact with someone of the opposite sex with known risk such as injection drug use, bisexual male (applies to females only), person with hemophilia/coagulation disorder, transfusion recipient with HIV documentation, and or person with AIDS or documented HIV. 2. Includes perinatal exposure current age ≥ 13; hemophilia, blood transfusion. 3. Perinatal/other pediatric: cases <13 born to a mother who is HIV-positive, and cases <13 with other risk (e.g., sexual transmission). 4. Adjusted for missing risk using multiple imputation methods. Adjusted subtotals may be different from unadjusted subtotals due to rounding.

TABLE 17. HIV DIAGNOSES AND PEOPLE WITH HIV (PWH), DISTRICT 2 (GAINESVILLE), GA, 2024

	HIV Diagnoses		PWH	
	N	%*	N	%*
Total	64	100.0	1,242	100.0
Birth Sex				
Males	44	68.8	892	71.8
Females	20	31.3	349	28.1
Gender Identity				
Cisgender Men	43	67.2	877	70.6
Cisgender Women	20	31.3	349	28.1
Transgender Persons/Additional Gender Identity	<5	---	15	1.2
Race and Ethnicity				
Black or African American, non-Hispanic	11	17.2	304	24.5
White, non-Hispanic	26	40.6	550	44.3
Hispanic or Latino, Any Race	22	34.4	255	20.5
Asian	0	0.0	21	1.7
American Indian or Alaska Native	0	0.0	<5	---
Native Hawaiian or Pacific Islander	0	0.0	0	0.0
Multiple Races	<5	---	89	7.2
Age Group (at diagnosis for diagnosed, current for PWH)				
<13	0	0.0	0	0.0
13-17	<5	---	<5	---
18-24	5	7.8	29	2.3
25-34	27	42.2	198	15.9
35-44	16	25.0	267	21.5
45-54	5	7.8	273	22.0
55-64	6	9.4	300	24.2
65+	<5	---	173	13.9
HIV Transmission Category (Unadjusted)				
Male-to-male sexual contact (MMSC)	28	43.8	542	43.6
Heterosexual contact ¹	22	34.4	237	19.1
Injection drug use (IDU)	6	9.4	67	5.4
MMSC and IDU	<5	---	60	4.8
Adult Other ²	<5	---	16	1.3
Perinatal/other pediatric ³	0	0.0	0	0.0
No risk identified or reported (NIR/NRR)	6	9.4	320	25.8
HIV Transmission Category (Adjusted)⁴				
Male-to-male sexual contact (MMSC)	32	49.5	682	54.9
Heterosexual contact ¹	23	36.1	360	29.0
Injection drug use (IDU)	6	9.5	90	7.2
MMSC and IDU	<5	---	72	5.8
Adult Other ²	<5	---	16	1.3
Perinatal/other pediatric ³	0	0.0	0	0.0
No risk identified or reported (NIR/NRR)	<5	---	22	1.8

* Percentages may not add up to 100% due to rounding 1. Heterosexual contact: Defined as sexual contact with someone of the opposite sex with known risk such as injection drug use, bisexual male (applies to females only), person with hemophilia/coagulation disorder, transfusion recipient with HIV documentation, and or person with AIDS or documented HIV. 2. Includes perinatal exposure current age \geq 13; hemophilia, blood transfusion. 3. Perinatal/other pediatric: cases <13 born to a mother who is HIV-positive, and cases <13 with other risk (e.g., sexual transmission). 4. Adjusted for missing risk using multiple imputation methods. Adjusted subtotals may be different from unadjusted subtotals due to rounding.

TABLE 18. HIV DIAGNOSES AND PEOPLE WITH HIV (PWH), DISTRICT 3-1 (COBB-DOUGLAS), GA, 2024

	HIV Diagnoses		PWH	
	N	%*	N	%*
Total	195	100.0	5,015	100.0
Birth Sex				
Males	152	77.9	3,935	78.5
Females	43	22.1	1,075	21.4
Gender Identity				
Cisgender Men	149	76.4	3,838	76.5
Cisgender Women	43	22.1	1,073	21.4
Transgender Persons/Additional Gender Identity	<5	---	99	2.0
Race and Ethnicity				
Black or African American, non-Hispanic	121	62.1	3,066	61.1
White, non-Hispanic	15	7.7	863	17.2
Hispanic or Latino, Any Race	55	28.2	691	13.8
Asian	0	0.0	20	0.4
American Indian or Alaska Native	<5	---	<5	---
Native Hawaiian or Pacific Islander	0	0.0	<5	---
Multiple Races	<5	---	333	6.6
Age Group (at diagnosis for diagnosed, current for PWH)				
<13	0	0.0	6	0.1
13-17	<5	---	<5	---
18-24	45	23.1	181	3.6
25-34	71	36.4	1,036	20.7
35-44	39	20.0	1,259	25.1
45-54	29	14.9	1,010	20.1
55-64	8	4.1	1,045	20.8
65+	<5	---	474	9.5
HIV Transmission Category (Unadjusted)				
Male-to-male sexual contact (MMSC)	117	60.0	2,892	57.7
Heterosexual contact ¹	59	30.3	908	18.1
Injection drug use (IDU)	<5	---	121	2.4
MMSC and IDU	<5	---	153	3.1
Adult Other ²	0	0.0	51	1.0
Perinatal/other pediatric ³	0	0.0	6	0.1
No risk identified or reported (NIR/NRR)	17	8.7	884	17.6
HIV Transmission Category (Adjusted)⁴				
Male-to-male sexual contact (MMSC)	129	66.2	3,335	66.5
Heterosexual contact ¹	62	31.8	1,222	24.4
Injection drug use (IDU)	<5	---	175	3.5
MMSC and IDU	<5	---	186	3.7
Adult Other ²	0	0.0	51	1.0
Perinatal/other pediatric ³	0	0.0	6	0.1
No risk identified or reported (NIR/NRR)	<5	---	41	0.8

* Percentages may not add up to 100% due to rounding. 1. Heterosexual contact: Defined as sexual contact with someone of the opposite sex with known risk such as injection drug use, bisexual male (applies to females only), person with hemophilia/coagulation disorder, transfusion recipient with HIV documentation, and or person with AIDS or documented HIV. 2. Includes perinatal exposure current age ≥ 13; hemophilia, blood transfusion. 3. Perinatal/other pediatric: cases <13 born to a mother who is HIV-positive, and cases <13 with other risk (e.g., sexual transmission). 4. Adjusted for missing risk using multiple imputation methods. Adjusted subtotals may be different from unadjusted subtotals due to rounding.

TABLE 19. HIV DIAGNOSES AND PEOPLE WITH HIV (PWH), DISTRICT 3-2 (FULTON), GA, 2024

	HIV Diagnoses		PWH	
	N	%*	N	%*
Total	481	100.0	17,441	100.0
Birth Sex				
Males	389	80.9	14,852	85.2
Females	92	19.1	2,574	14.8
Gender Identity				
Cisgender Men	374	77.8	14,512	83.2
Cisgender Women	91	18.9	2,564	14.7
Transgender Persons/Additional Gender Identity	16	3.3	350	2.0
Race and Ethnicity				
Black or African American, non-Hispanic	350	72.8	12,626	72.4
White, non-Hispanic	55	11.4	2,318	13.3
Hispanic or Latino, Any Race	52	10.8	1,218	7.0
Asian	5	1.0	62	0.4
American Indian or Alaska Native	<5	---	8	0.0
Native Hawaiian or Pacific Islander	0	0.0	<5	---
Multiple Races	16	3.3	1,044	6.0
Age Group (at diagnosis for diagnosed, current for PWH)				
<13	<5	---	12	0.1
13-17	<5	---	14	0.1
18-24	85	17.7	381	2.2
25-34	196	40.7	3,251	18.6
35-44	112	23.3	4,630	26.5
45-54	44	9.1	3,385	19.4
55-64	26	5.4	3,726	21.4
65+	14	2.9	2,041	11.7
HIV Transmission Category (Unadjusted)				
Male-to-male sexual contact (MMSC)	283	58.8	11,304	64.8
Heterosexual contact ¹	121	25.2	2,125	12.2
Injection drug use (IDU)	9	1.9	534	3.1
MMSC and IDU	7	1.5	642	3.7
Adult Other ²	0	0.0	109	0.6
Perinatal/other pediatric ³	<5	---	10	0.1
No risk identified or reported (NIR/NRR)	59	12.3	2,717	15.6
HIV Transmission Category (Adjusted)⁴				
Male-to-male sexual contact (MMSC)	318	66.2	12,748	73.1
Heterosexual contact ¹	140	29.1	2,986	17.1
Injection drug use (IDU)	10	2.2	690	4.0
MMSC and IDU	8	1.7	738	4.2
Adult Other ²	0	0.0	109	0.6
Perinatal/other pediatric ³	<5	---	10	0.1
No risk identified or reported (NIR/NRR)	<5	---	160	0.9

* Percentages may not add up to 100% due to rounding. 1. Heterosexual contact: Defined as sexual contact with someone of the opposite sex with known risk such as injection drug use, bisexual male (applies to females only), person with hemophilia/coagulation disorder, transfusion recipient with HIV documentation, and or person with AIDS or documented HIV. 2. Includes perinatal exposure current age ≥ 13; hemophilia, blood transfusion. 3. Perinatal/other pediatric: cases <13 born to a mother who is HIV-positive, and cases <13 with other risk (e.g., sexual transmission). 4. Adjusted for missing risk using multiple imputation methods. Adjusted subtotals may be different from unadjusted subtotals due to rounding.

TABLE 20. HIV DIAGNOSES AND PEOPLE WITH HIV (PWH), DISTRICT 3-3 (CLAYTON), GA, 2024

	HIV Diagnoses		PWH	
	N	%*	N	%*
Total	133	100.0	3,437	100.0
Birth Sex				
Males	104	78.2	2,643	76.9
Females	29	21.8	794	23.1
Gender Identity				
Cisgender Men	101	75.9	2,584	75.2
Cisgender Women	29	21.8	791	23.0
Transgender Persons/Additional Gender Identity	<5	---	62	1.8
Race and Ethnicity				
Black or African American, non-Hispanic	105	78.9	2,801	81.5
White, non-Hispanic	<5	---	110	3.2
Hispanic or Latino, Any Race	19	14.3	266	7.7
Asian	<5	---	26	0.8
American Indian or Alaska Native	0	0.0	<5	---
Native Hawaiian or Pacific Islander	0	0.0	<5	---
Multiple Races	<5	---	208	6.1
Age Group (at diagnosis for diagnosed, current for PWH)				
<13	<5	---	<5	---
13-17	<5	---	<5	---
18-24	38	28.6	117	3.4
25-34	59	44.4	813	23.7
35-44	13	9.8	906	26.4
45-54	9	6.8	639	18.6
55-64	6	4.5	653	19.0
65+	<5	---	301	8.8
HIV Transmission Category (Unadjusted)				
Male-to-male sexual contact (MMSC)	69	51.9	1,968	57.3
Heterosexual contact ¹	46	34.6	640	18.6
Injection drug use (IDU)	<5	---	101	2.9
MMSC and IDU	0	0.0	86	2.5
Adult Other ²	0	0.0	25	0.7
Perinatal/other pediatric ³	<5	---	<5	---
No risk identified or reported (NIR/NRR)	14	10.5	613	17.8
HIV Transmission Category (Adjusted)⁴				
Male-to-male sexual contact (MMSC)	75	56.0	2,232	65.0
Heterosexual contact ¹	53	39.7	910	26.5
Injection drug use (IDU)	<5	---	139	4.0
MMSC and IDU	<5	---	105	3.1
Adult Other ²	0	0.0	25	0.7
Perinatal/other pediatric ³	<5	---	<5	---
No risk identified or reported (NIR/NRR)	<5	---	22	0.6

* Percentages may not add up to 100% due to rounding. 1. Heterosexual contact: Defined as sexual contact with someone of the opposite sex with known risk such as injection drug use, bisexual male (applies to females only), person with hemophilia/coagulation disorder, transfusion recipient with HIV documentation, and or person with AIDS or documented HIV. 2. Includes perinatal exposure current age \geq 13; hemophilia, blood transfusion. 3. Perinatal/other pediatric: cases <13 born to a mother who is HIV-positive, and cases <13 with other risk (e.g., sexual transmission). 4. Adjusted for missing risk using multiple imputation methods. Adjusted subtotals may be different from unadjusted subtotals due to rounding.

TABLE 21. HIV DIAGNOSES AND PEOPLE WITH HIV (PWH), DISTRICT 3-4 (GNR), GA, 2024

	HIV Diagnoses		PWH	
	N	%*	N	%*
Total	267	100.0	5,529	100.0
Birth Sex				
Males	217	81.3	4,073	73.7
Females	50	18.7	1,451	26.2
Gender Identity				
Cisgender Men	212	79.4	3,984	72.1
Cisgender Women	50	18.7	1,444	26.1
Transgender Persons/Additional Gender Identity	5	1.9	96	1.7
Race and Ethnicity				
Black or African American, non-Hispanic	127	47.6	3,169	57.3
White, non-Hispanic	19	7.1	686	12.4
Hispanic or Latino, Any Race	104	39.0	1,192	21.6
Asian	10	3.7	110	2.0
American Indian or Alaska Native	0	0.0	<5	---
Native Hawaiian or Pacific Islander	0	0.0	<5	---
Multiple Races	7	2.6	311	5.6
Age Group (at diagnosis for diagnosed, current for PWH)				
<13	0	0.0	6	0.1
13-17	<5	---	8	0.1
18-24	45	16.9	199	3.6
25-34	105	39.3	1,104	20.0
35-44	58	21.7	1,350	24.4
45-54	26	9.7	1,184	21.4
55-64	22	8.2	1,155	20.9
65+	10	3.7	523	9.5
HIV Transmission Category (Unadjusted)				
Male-to-male sexual contact (MMSC)	157	58.8	2,877	52.0
Heterosexual contact ¹	89	33.3	1,077	19.5
Injection drug use (IDU)	<5	---	160	2.9
MMSC and IDU	<5	---	129	2.3
Adult Other ²	<5	---	64	1.2
Perinatal/other pediatric ³	0	0.0	6	0.1
No risk identified or reported (NIR/NRR)	17	6.4	1,216	22.0
HIV Transmission Category (Adjusted)⁴				
Male-to-male sexual contact (MMSC)	168	62.9	3,415	61.8
Heterosexual contact ¹	94	35.2	1,588	28.7
Injection drug use (IDU)	<5	---	234	4.2
MMSC and IDU	<5	---	166	3.0
Adult Other ²	<5	---	64	1.2
Perinatal/other pediatric ³	0	0.0	6	0.1
No risk identified or reported (NIR/NRR)	0	0.0	56	1.0

* Percentages may not add up to 100% due to rounding. 1. Heterosexual contact: Defined as sexual contact with someone of the opposite sex with known risk such as injection drug use, bisexual male (applies to females only), person with hemophilia/coagulation disorder, transfusion recipient with HIV documentation, and or person with AIDS or documented HIV. 2. Includes perinatal exposure current age ≥ 13 ; hemophilia, blood transfusion. 3. Perinatal/other pediatric: cases <13 born to a mother who is HIV-positive, and cases <13 with other risk (e.g., sexual transmission). 4. Adjusted for missing risk using multiple imputation methods. Adjusted subtotals may be different from unadjusted subtotals due to rounding.

TABLE 22. HIV DIAGNOSES AND PEOPLE WITH HIV (PWH), DISTRICT 3-5 (DEKALB), GA, 2024

	HIV Diagnoses		PWH	
	N	%*	N	%*
Total	319	100.0	10,302	100.0
Birth Sex				
Males	245	76.8	8,388	81.4
Females	74	23.2	1,906	18.5
Gender Identity				
Cisgender Men	237	74.3	8,196	79.6
Cisgender Women	74	23.2	1,899	18.4
Transgender Persons/Additional Gender Identity	8	2.5	199	1.9
Race and Ethnicity				
Black or African American, non-Hispanic	223	69.9	7,271	70.6
White, non-Hispanic	31	9.7	1,346	13.1
Hispanic or Latino, Any Race	48	15.0	915	8.9
Asian	7	2.2	80	0.8
American Indian or Alaska Native	<5	---	<5	---
Native Hawaiian or Pacific Islander	0	0.0	<5	---
Multiple Races	7	2.2	602	5.8
Age Group (at diagnosis for diagnosed, current for PWH)				
<13	0	0.0	15	0.1
13-17	<5	---	12	0.1
18-24	56	17.6	254	2.5
25-34	125	39.2	1,825	17.7
35-44	79	24.8	2,553	24.8
45-54	30	9.4	2,078	20.2
55-64	19	6.0	2,378	23.1
65+	7	2.2	1,187	11.5
HIV Transmission Category (Unadjusted)				
Male-to-male sexual contact (MMSC)	179	56.1	6,307	61.2
Heterosexual contact ¹	84	26.3	1,517	14.7
Injection drug use (IDU)	<5	---	239	2.3
MMSC and IDU	<5	---	267	2.6
Adult Other ²	<5	---	90	0.9
Perinatal/other pediatric ³	0	0.0	13	0.1
No risk identified or reported (NIR/NRR)	52	16.3	1,869	18.1
HIV Transmission Category (Adjusted)⁴				
Male-to-male sexual contact (MMSC)	203	63.5	7,237	70.3
Heterosexual contact ¹	107	33.4	2,174	21.1
Injection drug use (IDU)	5	1.6	365	3.5
MMSC and IDU	<5	---	337	3.3
Adult Other ²	<5	---	90	0.9
Perinatal/other pediatric ³	0	0.0	13	0.1
No risk identified or reported (NIR/NRR)	<5	---	85	0.8

* Percentages may not add up to 100% due to rounding. 1. Heterosexual contact: Defined as sexual contact with someone of the opposite sex with known risk such as injection drug use, bisexual male (applies to females only), person with hemophilia/coagulation disorder, transfusion recipient with HIV documentation, and or person with AIDS or documented HIV. 2. Includes perinatal exposure current age ≥ 13; hemophilia, blood transfusion. 3. Perinatal/other pediatric: cases <13 born to a mother who is HIV-positive, and cases <13 with other risk (e.g., sexual transmission). 4. Adjusted for missing risk using multiple imputation methods. Adjusted subtotals may be different from unadjusted subtotals due to rounding.

TABLE 23. HIV DIAGNOSES AND PEOPLE WITH HIV (PWH), DISTRICT 4 (LAGRANGE), GA, 2024

	HIV Diagnoses		PWH	
	N	%*	N	%*
Total	160	100.0	3,225	100.0
Birth Sex				
Males	128	80.0	2,359	73.1
Females	32	20.0	864	26.8
Gender Identity				
Cisgender Men	123	76.9	2,316	71.8
Cisgender Women	32	20.0	862	26.7
Transgender Persons/Additional Gender Identity	5	3.1	45	1.4
Race and Ethnicity				
Black or African American, non-Hispanic	114	71.3	2,163	67.1
White, non-Hispanic	26	16.3	579	18.0
Hispanic or Latino, Any Race	15	9.4	252	7.8
Asian	<5	---	11	0.3
American Indian or Alaska Native	0	0.0	<5	---
Native Hawaiian or Pacific Islander	0	0.0	0	0.0
Multiple Races	<5	---	193	6.0
Age Group (at diagnosis for diagnosed, current for PWH)				
<13	<5	---	6	0.2
13-17	<5	---	8	0.2
18-24	42	26.3	148	4.6
25-34	55	34.4	582	18.0
35-44	28	17.5	749	23.2
45-54	21	13.1	662	20.5
55-64	5	3.1	661	20.5
65+	<5	---	409	12.7
HIV Transmission Category (Unadjusted)				
Male-to-male sexual contact (MMSC)	80	50.0	1,529	47.4
Heterosexual contact ¹	56	35.0	696	21.6
Injection drug use (IDU)	<5	---	113	3.5
MMSC and IDU	<5	---	104	3.2
Adult Other ²	0	0.0	36	1.1
Perinatal/other pediatric ³	<5	---	5	0.2
No risk identified or reported (NIR/NRR)	20	12.5	742	23.0
HIV Transmission Category (Adjusted)⁴				
Male-to-male sexual contact (MMSC)	95	59.3	1,875	58.1
Heterosexual contact ¹	60	37.5	991	30.7
Injection drug use (IDU)	<5	---	165	5.1
MMSC and IDU	<5	---	127	4.0
Adult Other ²	0	0.0	36	1.1
Perinatal/other pediatric ³	<5	---	5	0.2
No risk identified or reported (NIR/NRR)	0	0.0	26	0.8

* Percentages may not add up to 100% due to rounding 1. Heterosexual contact: Defined as sexual contact with someone of the opposite sex with known risk such as injection drug use, bisexual male (applies to females only), person with hemophilia/coagulation disorder, transfusion recipient with HIV documentation, and or person with AIDS or documented HIV. 2. Includes perinatal exposure current age \geq 13; hemophilia, blood transfusion. 3. Perinatal/other pediatric: cases <13 born to a mother who is HIV-positive, and cases <13 with other risk (e.g., sexual transmission). 4. Adjusted for missing risk using multiple imputation methods. Adjusted subtotals may be different from unadjusted subtotals due to rounding.

TABLE 24. HIV DIAGNOSES AND PEOPLE WITH HIV (PWH), DISTRICT 5-1 (DUBLIN), GA, 2024

	HIV Diagnoses		PWH	
	N	%*	N	%*
Total	24	100.0	731	100.0
Birth Sex				
Males	17	70.8	507	69.4
Females	7	29.2	224	30.6
Gender Identity				
Cisgender Men	17	70.8	494	67.6
Cisgender Women	7	29.2	224	30.6
Transgender Persons/Additional Gender Identity	0	0.0	13	1.8
Race and Ethnicity				
Black or African American, non-Hispanic	18	75.0	499	68.3
White, non-Hispanic	<5	---	92	12.6
Hispanic or Latino, Any Race	<5	---	64	8.8
Asian	0	0.0	0	0.0
American Indian or Alaska Native	0	0.0	0	0.0
Native Hawaiian or Pacific Islander	0	0.0	0	0.0
Multiple Races	0	0.0	70	9.6
Age Group (at diagnosis for diagnosed, current for PWH)				
<13	0	0.0	<5	---
13-17	0	0.0	<5	---
18-24	6	25.0	23	3.1
25-34	10	41.7	94	12.9
35-44	<5	---	154	21.1
45-54	<5	---	167	22.8
55-64	<5	---	194	26.5
65+	0	0.0	96	13.1
HIV Transmission Category (Unadjusted)				
Male-to-male sexual contact (MMSC)	10	41.7	223	30.5
Heterosexual contact ¹	7	29.2	171	23.4
Injection drug use (IDU)	<5	---	46	6.3
MMSC and IDU	0	0.0	31	4.2
Adult Other ²	0	0.0	7	1.0
Perinatal/other pediatric ³	0	0.0	<5	---
No risk identified or reported (NIR/NRR)	6	25.0	251	34.3
HIV Transmission Category (Adjusted)⁴				
Male-to-male sexual contact (MMSC)	14	60.0	345	47.3
Heterosexual contact ¹	9	35.8	265	36.2
Injection drug use (IDU)	<5	---	67	9.1
MMSC and IDU	0	0.0	39	5.4
Adult Other ²	0	0.0	7	1.0
Perinatal/other pediatric ³	0	0.0	<5	---
No risk identified or reported (NIR/NRR)	0	0.0	6	0.8

* Percentages may not add up to 100% due to rounding. 1. Heterosexual contact: Defined as sexual contact with someone of the opposite sex with known risk such as injection drug use, bisexual male (applies to females only), person with hemophilia/coagulation disorder, transfusion recipient with HIV documentation, and or person with AIDS or documented HIV. 2. Includes perinatal exposure current age \geq 13; hemophilia, blood transfusion. 3. Perinatal/other pediatric: cases <13 born to a mother who is HIV-positive, and cases <13 with other risk (e.g., sexual transmission). 4. Adjusted for missing risk using multiple imputation methods. Adjusted subtotals may be different from unadjusted subtotals due to rounding.

TABLE 25. HIV DIAGNOSES AND PEOPLE WITH HIV (PWH), DISTRICT 5-2 (MACON), GA, 2024

	HIV Diagnoses		PWH	
	N	%*	N	%*
Total	126	100.0	2,803	100.0
Birth Sex				
Males	83	65.9	1,940	69.2
Females	43	34.1	862	30.8
Gender Identity				
Cisgender Men	82	65.1	1,904	67.9
Cisgender Women	43	34.1	860	30.7
Transgender Persons/Additional Gender Identity	<5	---	38	1.4
Race and Ethnicity				
Black or African American, non-Hispanic	87	69.0	2,062	73.6
White, non-Hispanic	24	19.0	392	14.0
Hispanic or Latino, Any Race	10	7.9	145	5.2
Asian	0	0.0	<5	---
American Indian or Alaska Native	<5	---	<5	---
Native Hawaiian or Pacific Islander	0	0.0	0	0.0
Multiple Races	<5	---	181	6.5
Age Group (at diagnosis for diagnosed, current for PWH)				
<13	0	0.0	<5	---
13-17	<5	---	<5	---
18-24	19	15.1	96	3.4
25-34	53	42.1	531	18.9
35-44	29	23.0	645	23.0
45-54	10	7.9	533	19.0
55-64	6	4.8	637	22.7
65+	7	5.6	354	12.6
HIV Transmission Category (Unadjusted)				
Male-to-male sexual contact (MMSC)	56	44.4	1,166	41.6
Heterosexual contact ¹	54	42.9	646	23.0
Injection drug use (IDU)	<5	---	116	4.1
MMSC and IDU	<5	---	93	3.3
Adult Other ²	0	0.0	26	0.9
Perinatal/other pediatric ³	0	0.0	<5	---
No risk identified or reported (NIR/NRR)	14	11.1	754	26.9
HIV Transmission Category (Adjusted)⁴				
Male-to-male sexual contact (MMSC)	66	52.2	1,486	53.0
Heterosexual contact ¹	58	45.6	987	35.2
Injection drug use (IDU)	<5	---	172	6.1
MMSC and IDU	<5	---	112	4.0
Adult Other ²	0	0.0	26	0.9
Perinatal/other pediatric ³	0	0.0	<5	---
No risk identified or reported (NIR/NRR)	0	0.0	18	0.6

* Percentages may not add up to 100% due to rounding 1. Heterosexual contact: Defined as sexual contact with someone of the opposite sex with known risk such as injection drug use, bisexual male (applies to females only), person with hemophilia/coagulation disorder, transfusion recipient with HIV documentation, and or person with AIDS or documented HIV. 2. Includes perinatal exposure current age \geq 13; hemophilia, blood transfusion. 3. Perinatal/other pediatric: cases <13 born to a mother who is HIV-positive, and cases <13 with other risk (e.g., sexual transmission). 4. Adjusted for missing risk using multiple imputation methods. Adjusted subtotals may be different from unadjusted subtotals due to rounding.

TABLE 26. HIV DIAGNOSES AND PEOPLE WITH HIV (PWH), DISTRICT 6 (AUGUSTA), GA, 2024

	HIV Diagnoses		PWH	
	N	%*	N	%*
Total	99	100	2,519	100
Birth Sex				
Males	83	83.8	1,762	69.9
Females	16	16.2	757	30.1
Gender Identity				
Cisgender Men	82	82.8	1,713	68
Cisgender Women	16	16.2	753	29.9
Transgender Persons/Additional Gender Identity	<5	---	53	2.1
Race and Ethnicity				
Black or African American, non-Hispanic	66	66.7	1,851	73.5
White, non-Hispanic	19	19.2	421	16.7
Hispanic or Latino, Any Race	10	10.1	109	4.3
Asian	<5	---	6	0.2
American Indian or Alaska Native	0	0	0	0
Native Hawaiian or Pacific Islander	0	0	<5	---
Multiple Races	<5	---	116	4.6
Age Group (at diagnosis for diagnosed, current for PWH)				
<13	<5	---	<5	---
13-17	0	0	<5	---
18-24	24	24.2	114	4.5
25-34	42	42.4	485	19.3
35-44	16	16.2	475	18.9
45-54	10	10.1	398	15.8
55-64	<5	---	617	24.5
65+	<5	---	423	16.8
HIV Transmission Category (Unadjusted)				
Male-to-male sexual contact (MMSC)	50	50.5	1,072	42.6
Heterosexual contact ¹	35	35.4	571	22.7
Injection drug use (IDU)	<5	---	142	5.6
MMSC and IDU	<5	---	94	3.7
Adult Other ²	0	0	80	3.2
Perinatal/other pediatric ³	<5	---	<5	---
No risk identified or reported (NIR/NRR)	11	11.1	556	22.1
HIV Transmission Category (Adjusted)⁴				
Male-to-male sexual contact (MMSC)	59	59.2	1,282	50.9
Heterosexual contact ¹	36	36	836	33.2
Injection drug use (IDU)	<5	---	194	7.7
MMSC and IDU	<5	---	107	4.3
Adult Other ²	0	0	80	3.2
Perinatal/other pediatric ³	<5	---	<5	---
No risk identified or reported (NIR/NRR)	<5	---	15	0.6

* Percentages may not add up to 100% due to rounding 1. Heterosexual contact: Defined as sexual contact with someone of the opposite sex with known risk such as injection drug use, bisexual male (applies to females only), person with hemophilia/coagulation disorder, transfusion recipient with HIV documentation, and or person with AIDS or documented HIV. 2. Includes perinatal exposure current age \geq 13; hemophilia, blood transfusion. 3. Perinatal/other pediatric: cases <13 born to a mother who is HIV-positive, and cases <13 with other risk (e.g., sexual transmission). 4. Adjusted for missing risk using multiple imputation methods. Adjusted subtotals may be different from unadjusted subtotals due to rounding.

TABLE 27. HIV DIAGNOSES AND PEOPLE WITH HIV (PWH), DISTRICT 7 (COLUMBUS), GA, 2024

	HIV Diagnoses		PWH	
	N	%*	N	%*
Total	91	100.0	2,115	100.0
Birth Sex				
Males	62	68.1	1,427	67.5
Females	29	31.9	687	32.5
Gender Identity				
Cisgender Men	61	67.0	1,407	66.5
Cisgender Women	29	31.9	686	32.4
Transgender Persons/Additional Gender Identity	<5	---	21	1.0
Race and Ethnicity				
Black or African American, non-Hispanic	71	78.0	1,526	72.2
White, non-Hispanic	13	14.3	250	11.8
Hispanic or Latino, Any Race	6	6.6	170	8.0
Asian	0	0.0	<5	---
American Indian or Alaska Native	0	0.0	<5	---
Native Hawaiian or Pacific Islander	0	0.0	0	0.0
Multiple Races	<5	---	124	5.9
Age Group (at diagnosis for diagnosed, current for PWH)				
<13	0	0.0	<5	---
13-17	<5	---	6	0.3
18-24	16	17.6	68	3.2
25-34	28	30.8	358	16.9
35-44	22	24.2	461	21.8
45-54	15	16.5	454	21.5
55-64	7	7.7	482	22.8
65+	<5	---	282	13.3
HIV Transmission Category (Unadjusted)				
Male-to-male sexual contact (MMSC)	33	36.3	745	35.2
Heterosexual contact ¹	40	44.0	609	28.8
Injection drug use (IDU)	<5	---	101	4.8
MMSC and IDU	<5	---	49	2.3
Adult Other ²	0	0.0	23	1.1
Perinatal/other pediatric ³	0	0.0	<5	---
No risk identified or reported (NIR/NRR)	15	16.5	584	27.6
HIV Transmission Category (Adjusted)⁴				
Male-to-male sexual contact (MMSC)	43	46.8	1,001	47.3
Heterosexual contact ¹	44	48.8	841	39.8
Injection drug use (IDU)	<5	---	139	6.6
MMSC and IDU	<5	---	64	3.0
Adult Other ²	0	0.0	23	1.1
Perinatal/other pediatric ³	0	0.0	<5	---
No risk identified or reported (NIR/NRR)	0	0.0	43	2.0

* Percentages may not add up to 100% due to rounding 1. Heterosexual contact: Defined as sexual contact with someone of the opposite sex with known risk such as injection drug use, bisexual male (applies to females only), person with hemophilia/coagulation disorder, transfusion recipient with HIV documentation, and or person with AIDS or documented HIV. 2. Includes perinatal exposure current age ≥ 13; hemophilia, blood transfusion. 3. Perinatal/other pediatric: cases <13 born to a mother who is HIV-positive, and cases <13 with other risk (e.g., sexual transmission). 4. Adjusted for missing risk using multiple imputation methods. Adjusted subtotals may be different from unadjusted subtotals due to rounding.

TABLE 28. HIV DIAGNOSES AND PEOPLE WITH HIV (PWH), DISTRICT 8-1 (VALDOSTA), GA, 2024

	HIV Diagnoses		PWH	
	N	%*	N	%*
Total	62	100.0	1,362	100.0
Birth Sex				
Males	38	61.3	844	62.0
Females	24	38.7	518	38.0
Gender Identity				
Cisgender Men	37	59.7	831	61.0
Cisgender Women	24	38.7	517	38.0
Transgender Persons/Additional Gender Identity	<5	---	14	1.0
Race and Ethnicity				
Black or African American, non-Hispanic	42	67.7	923	67.8
White, non-Hispanic	15	24.2	230	16.9
Hispanic or Latino, Any Race	<5	---	116	8.5
Asian	0	0.0	<5	---
American Indian or Alaska Native	0	0.0	0	0.0
Native Hawaiian or Pacific Islander	0	0.0	0	0.0
Multiple Races	0	0.0	83	6.1
Age Group (at diagnosis for diagnosed, current for PWH)				
<13	0	0.0	<5	---
13-17	0	0.0	<5	---
18-24	9	14.5	41	3.0
25-34	23	37.1	200	14.7
35-44	12	19.4	285	20.9
45-54	8	12.9	294	21.6
55-64	8	12.9	325	23.9
65+	<5	---	215	15.8
HIV Transmission Category (Unadjusted)				
Male-to-male sexual contact (MMSC)	21	33.9	433	31.8
Heterosexual contact ¹	23	37.1	498	36.6
Injection drug use (IDU)	<5	---	58	4.3
MMSC and IDU	<5	---	36	2.6
Adult Other ²	0	0.0	12	0.9
Perinatal/other pediatric ³	0	0.0	<5	---
No risk identified or reported (NIR/NRR)	16	25.8	324	23.8
HIV Transmission Category (Adjusted)⁴				
Male-to-male sexual contact (MMSC)	28	45.0	558	41.0
Heterosexual contact ¹	31	49.7	658	48.3
Injection drug use (IDU)	<5	---	83	6.1
MMSC and IDU	<5	---	43	3.2
Adult Other ²	0	0.0	12	0.9
Perinatal/other pediatric ³	0	0.0	<5	---
No risk identified or reported (NIR/NRR)	0	0.0	6	0.4

* Percentages may not add up to 100% due to rounding. 1. Heterosexual contact: Defined as sexual contact with someone of the opposite sex with known risk such as injection drug use, bisexual male (applies to females only), person with hemophilia/coagulation disorder, transfusion recipient with HIV documentation, and or person with AIDS or documented HIV. 2. Includes perinatal exposure current age \geq 13; hemophilia, blood transfusion. 3. Perinatal/other pediatric: cases <13 born to a mother who is HIV-positive, and cases <13 with other risk (e.g., sexual transmission). 4. Adjusted for missing risk using multiple imputation methods. Adjusted subtotals may be different from unadjusted subtotals due to rounding.

TABLE 29. HIV DIAGNOSES AND PEOPLE WITH HIV (PWH), DISTRICT 8-2 (ALBANY), GA, 2024

	HIV Diagnoses		PWH	
	N	%*	N	%*
Total	70	100.0	1,948	100.0
Birth Sex				
Males	54	77.1	1,258	64.6
Females	16	22.9	690	35.4
Gender Identity				
Cisgender Men	54	77.1	1,234	63.3
Cisgender Women	16	22.9	689	35.4
Transgender Persons/Additional Gender Identity	0	0.0	25	1.3
Race and Ethnicity				
Black or African American, non-Hispanic	49	70.0	1,505	77.3
White, non-Hispanic	12	17.1	211	10.8
Hispanic or Latino, Any Race	5	7.1	104	5.3
Asian	0	0.0	<5	---
American Indian or Alaska Native	0	0.0	0	0.0
Native Hawaiian or Pacific Islander	0	0.0	0	0.0
Multiple Races	<5	---	114	5.9
Age Group (at diagnosis for diagnosed, current for PWH)				
<13	<5	---	<5	---
13-17	0	0.0	<5	---
18-24	20	28.6	65	3.3
25-34	18	25.7	326	16.7
35-44	18	25.7	439	22.5
45-54	11	15.7	390	20.0
55-64	<5	---	439	22.5
65+	<5	---	283	14.5
HIV Transmission Category (Unadjusted)				
Male-to-male sexual contact (MMSC)	34	48.6	702	36.0
Heterosexual contact ¹	21	30.0	509	26.1
Injection drug use (IDU)	<5	---	75	3.9
MMSC and IDU	<5	---	41	2.1
Adult Other ²	0	0.0	21	1.1
Perinatal/other pediatric ³	<5	---	<5	---
No risk identified or reported (NIR/NRR)	11	15.7	597	30.6
HIV Transmission Category (Adjusted)⁴				
Male-to-male sexual contact (MMSC)	39	56.0	912	46.8
Heterosexual contact ¹	26	36.4	823	42.3
Injection drug use (IDU)	<5	---	123	6.3
MMSC and IDU	<5	---	54	2.8
Adult Other ²	0	0.0	21	1.1
Perinatal/other pediatric ³	<5	---	<5	---
No risk identified or reported (NIR/NRR)	0	0.0	11	0.6

* Percentages may not add up to 100% due to rounding 1. Heterosexual contact: Defined as sexual contact with someone of the opposite sex with known risk such as injection drug use, bisexual male (applies to females only), person with hemophilia/coagulation disorder, transfusion recipient with HIV documentation, and or person with AIDS or documented HIV. 2. Includes perinatal exposure current age \geq 13; hemophilia, blood transfusion. 3. Perinatal/other pediatric: cases <13 born to a mother who is HIV-positive, and cases <13 with other risk (e.g., sexual transmission). 4. Adjusted for missing risk using multiple imputation methods. Adjusted subtotals may be different from unadjusted subtotals due to rounding.

TABLE 30. HIV DIAGNOSES AND PEOPLE WITH HIV (PWH), DISTRICT 9-1 (SAVANNAH), GA, 2024

	HIV Diagnoses		PWH	
	N	%*	N	%*
Total	148	100.0	3,128	100.0
Birth Sex				
Males	122	82.4	2,222	71.0
Females	25	16.9	904	28.9
Gender Identity				
Cisgender Men	116	78.4	2,175	69.5
Cisgender Women	25	16.9	904	28.9
Transgender Persons/Additional Gender Identity	6	4.1	47	1.5
Race and Ethnicity				
Black or African American, non-Hispanic	87	58.8	1,952	62.4
White, non-Hispanic	26	17.6	615	19.7
Hispanic or Latino, Any Race	23	15.5	282	9.0
Asian	<5	---	10	0.3
American Indian or Alaska Native	0	0.0	0	0.0
Native Hawaiian or Pacific Islander	0	0.0	0	0.0
Multiple Races	8	5.4	231	7.4
Age Group (at diagnosis for diagnosed, current for PWH)				
<13	0	0.0	<5	---
13-17	<5	---	8	0.3
18-24	40	27.0	122	3.9
25-34	37	25.0	534	17.1
35-44	37	25.0	706	22.6
45-54	13	8.8	559	17.9
55-64	12	8.1	726	23.2
65+	7	4.7	469	15.0
HIV Transmission Category (Unadjusted)				
Male-to-male sexual contact (MMSC)	60	40.5	1,337	42.7
Heterosexual contact ¹	26	17.6	625	20.0
Injection drug use (IDU)	<5	---	140	4.5
MMSC and IDU	<5	---	97	3.1
Adult Other ²	0	0.0	41	1.3
Perinatal/other pediatric ³	0	0.0	<5	---
No risk identified or reported (NIR/NRR)	60	40.5	885	28.3
HIV Transmission Category (Adjusted)⁴				
Male-to-male sexual contact (MMSC)	94	63.8	1,688	53.9
Heterosexual contact ¹	45	30.5	1,027	32.8
Injection drug use (IDU)	<5	---	209	6.7
MMSC and IDU	<5	---	121	3.9
Adult Other ²	0	0.0	41	1.3
Perinatal/other pediatric ³	0	0.0	<5	---
No risk identified or reported (NIR/NRR)	<5	---	40	1.3

* Percentages may not add up to 100% due to rounding 1. Heterosexual contact: Defined as sexual contact with someone of the opposite sex with known risk such as injection drug use, bisexual male (applies to females only), person with hemophilia/coagulation disorder, transfusion recipient with HIV documentation, and or person with AIDS or documented HIV. 2. Includes perinatal exposure current age ≥ 13; hemophilia, blood transfusion. 3. Perinatal/other pediatric: cases <13 born to a mother who is HIV-positive, and cases <13 with other risk (e.g., sexual transmission). 4. Adjusted for missing risk using multiple imputation methods. Adjusted subtotals may be different from unadjusted subtotals due to rounding.

TABLE 31. HIV DIAGNOSES AND PEOPLE WITH HIV (PWH), DISTRICT 9-2 (WAYCROSS), GA, 2024

	HIV Diagnoses		PWH	
	N	%*	N	%*
Total	66	100.0	1,491	100.0
Birth Sex				
Males	51	77.3	1,064	71.4
Females	15	22.7	427	28.6
Gender Identity				
Cisgender Men	46	69.7	1,032	69.2
Cisgender Women	15	22.7	425	28.5
Transgender Persons/Additional Gender Identity	5	7.6	34	2.3
Race and Ethnicity				
Black or African American, non-Hispanic	33	50.0	845	56.7
White, non-Hispanic	18	27.3	309	20.7
Hispanic or Latino, Any Race	12	18.2	194	13.0
Asian	0	0.0	<5	---
American Indian or Alaska Native	0	0.0	<5	---
Native Hawaiian or Pacific Islander	0	0.0	0	0.0
Multiple Races	<5	---	110	7.4
Age Group (at diagnosis for diagnosed, current for PWH)				
<13	0	0.0	0	0.0
13-17	<5	---	0	0.0
18-24	17	25.8	60	4.0
25-34	20	30.3	277	18.6
35-44	8	12.1	330	22.1
45-54	15	22.7	277	18.6
55-64	<5	---	363	24.3
65+	<5	---	184	12.3
HIV Transmission Category (Unadjusted)				
Male-to-male sexual contact (MMSC)	36	54.5	595	39.9
Heterosexual contact ¹	19	28.8	348	23.3
Injection drug use (IDU)	<5	---	67	4.5
MMSC and IDU	<5	---	58	3.9
Adult Other ²	0	0.0	16	1.1
Perinatal/other pediatric ³	0	0.0	0	0.0
No risk identified or reported (NIR/NRR)	5	7.6	407	27.3
HIV Transmission Category (Adjusted)⁴				
Male-to-male sexual contact (MMSC)	39	58.5	756	50.7
Heterosexual contact ¹	21	32.3	526	35.3
Injection drug use (IDU)	<5	---	97	6.5
MMSC and IDU	<5	---	70	4.7
Adult Other ²	0	0.0	16	1.1
Perinatal/other pediatric ³	0	0.0	0	0.0
No risk identified or reported (NIR/NRR)	0	0.0	26	1.7

* Percentages may not add up to 100% due to rounding 1. Heterosexual contact: Defined as sexual contact with someone of the opposite sex with known risk such as injection drug use, bisexual male (applies to females only), person with hemophilia/coagulation disorder, transfusion recipient with HIV documentation, and or person with AIDS or documented HIV. 2. Includes perinatal exposure current age ≥ 13; hemophilia, blood transfusion. 3. Perinatal/other pediatric: cases <13 born to a mother who is HIV-positive, and cases <13 with other risk (e.g., sexual transmission). 4. Adjusted for missing risk using multiple imputation methods. Adjusted subtotals may be different from unadjusted subtotals due to rounding.

TABLE 32. HIV DIAGNOSES AND PEOPLE WITH HIV (PWH), DISTRICT 10 (ATHENS), GA, 2024

	HIV Diagnoses		PWH	
	N	%*	N	%*
Total	78	100.0	1,341	100.0
Birth Sex				
Males	60	76.9	929	69.3
Females	18	23.1	412	30.7
Gender Identity				
Cisgender Men	57	73.1	903	67.3
Cisgender Women	18	23.1	410	30.6
Transgender Persons/Additional Gender Identity	<5	---	28	2.1
Race and Ethnicity				
Black or African American, non-Hispanic	30	38.5	713	53.2
White, non-Hispanic	27	34.6	362	27.0
Hispanic or Latino, Any Race	15	19.2	174	13.0
Asian	<5	---	10	0.7
American Indian or Alaska Native	0	0.0	0	0.0
Native Hawaiian or Pacific Islander	0	0.0	0	0.0
Multiple Races	<5	---	70	5.2
Age Group (at diagnosis for diagnosed, current for PWH)				
<13	<5	---	<5	---
13-17	0	0.0	<5	---
18-24	19	24.4	61	4.5
25-34	28	35.9	256	19.1
35-44	9	11.5	292	21.8
45-54	10	12.8	262	19.5
55-64	10	12.8	299	22.3
65+	<5	---	167	12.5
HIV Transmission Category (Unadjusted)				
Male-to-male sexual contact (MMSC)	46	59.0	596	44.4
Heterosexual contact ¹	22	28.2	307	22.9
Injection drug use (IDU)	<5	---	65	4.8
MMSC and IDU	<5	---	38	2.8
Adult Other ²	0	0.0	16	1.2
Perinatal/other pediatric ³	<5	---	<5	---
No risk identified or reported (NIR/NRR)	5	6.4	317	23.6
HIV Transmission Category (Adjusted)⁴				
Male-to-male sexual contact (MMSC)	50	64.0	728	54.3
Heterosexual contact ¹	23	29.1	450	33.5
Injection drug use (IDU)	<5	---	88	6.6
MMSC and IDU	<5	---	47	3.5
Adult Other ²	0	0.0	16	1.2
Perinatal/other pediatric ³	<5	---	<5	---
No risk identified or reported (NIR/NRR)	0	0.0	11	0.8

* Percentages may not add up to 100% due to rounding 1. Heterosexual contact: Defined as sexual contact with someone of the opposite sex with known risk such as injection drug use, bisexual male (applies to females only), person with hemophilia/coagulation disorder, transfusion recipient with HIV documentation, and or person with AIDS or documented HIV. 2. Includes perinatal exposure current age ≥ 13; hemophilia, blood transfusion. 3.

Perinatal/other pediatric: cases <13 born to a mother who is HIV-positive, and cases <13 with other risk (e.g., sexual transmission). 4. Adjusted for missing risk using multiple imputation methods. Adjusted subtotals may be different from unadjusted subtotals due to rounding.

HIV DIAGNOSES AND PEOPLE WITH HIV, ATLANTA MSA

TABLE 33. HIV DIAGNOSES AND PEOPLE WITH HIV, ATLANTA MSA¹, GA, 2024

	HIV Diagnoses		PWH	
	N	%*	N	%*
Total	1,632	100.0	46,912	100.0
Birth Sex				
Males	1,284	78.7	37,658	80.3
Females	348	21.3	9,219	19.7
Gender Identity				
Cisgender Men	1,244	76.2	36,813	78.5
Cisgender Women	347	21.3	9,185	19.6
Transgender Persons/Additional Gender Identity	41	2.5	879	1.9
Race and Ethnicity				
Black or African American, non-Hispanic	1,063	65.1	31,859	67.9
White, non-Hispanic	183	11.2	6,634	14.1
Hispanic or Latino, Any Race	306	18.8	4,829	10.3
Asian	26	1.6	336	0.7
American Indian or Alaska Native	5	0.3	19	0.0
Native Hawaiian or Pacific Islander	0	0.0	9	0.0
Multiple Races	42	2.6	2,811	6.0
Age Group (at diagnosis for diagnosed, current for PWH)				
<13	5	0.3	52	0.1
13-17	14	0.9	53	0.1
18-24	323	19.8	1,322	2.8
25-34	638	39.1	8,951	19.1
35-44	348	21.3	11,864	25.3
45-54	163	10.0	9,397	20.0
55-64	96	5.9	10,101	21.5
65+	45	2.8	5,171	11.0
HIV Transmission Category (Unadjusted)				
Male-to-male sexual contact (MMSC)	911	55.8	27,745	59.1
Heterosexual contact ²	494	30.3	7,403	15.8
Injection drug use (IDU)	21	1.3	1,352	2.9
MMSC and IDU	16	1.0	1,456	3.1
Adult Other ³	<5	---	406	0.9
Perinatal/other pediatric ⁴	5	0.3	47	0.1
No risk identified or reported (NIR/NRR)	183	11.2	8,503	18.1
HIV Transmission Category (Adjusted)⁵				
Male-to-male sexual contact (MMSC)	1,016	62.3	31,929	68.1
Heterosexual contact ²	556	34.0	10,482	22.3
Injection drug use (IDU)	28	1.7	1,883	4.0
MMSC and IDU	19	1.2	1,751	3.7
Adult Other ³	<5	---	406	0.9
Perinatal/other pediatric ⁴	5	0.3	47	0.1
No risk identified or reported (NIR/NRR)	6	0.4	414	0.9

*Percents may not add up to 100 due to rounding. 1. Atlanta Metropolitan Statistical Area (MSA) counties consist of: Barrow, Bartow, Butts, Carroll, Cherokee, Clayton, Cobb, Coweta, Dawson, DeKalb, Douglas, Fayette, Forsyth, Fulton, Gwinnett, Haralson, Heard, Henry, Jasper, Lamar, Meriwether, Morgan, Newton, Paulding, Pickens, Pike, Rockdale, Spalding, Walton. 2. Heterosexual contact: Defined as sexual contact with someone of the opposite sex with known risk such as injection drug use, bisexual male (applies to females only), person with hemophilia/coagulation disorder, transfusion recipient with HIV documentation, and or person with AIDS or documented HIV. 3. Includes perinatal exposure age>13, hemophilia, blood transfusion. 4. Perinatal/other pediatric: cases <13 years born to a mother who is HIV-positive, and cases <13 with other risk (e.g., sexual transmission). 5. Adjusted for missing risk using multiple imputation methods. Adjusted subtotals may be different from unadjusted subtotals due to rounding.

HIV DIAGNOSES AND LATE DIAGNOSES BY PUBLIC HEALTH DISTRICT AND ATLANTA MSA

TABLE 34. HIV DIAGNOSES AND LATE DIAGNOSES BY PUBLIC HEALTH DISTRICT AND ATLANTA MSA, GA, 2024

	HIV Diagnoses		Late HIV Diagnoses ¹		Row %
	N	%*	N	%*	
Public Health District					
1-1 Northwest (Rome)	77	3.1	28	5.3	36.4
1-2 North Georgia (Dalton)	19	0.8	7	1.3	36.8
2 North (Gainesville)	64	2.6	19	3.6	29.7
3-1 Cobb-Douglas	195	7.8	40	7.5	20.5
3-2 Fulton	481	19.4	66	12.4	13.7
3-3 Clayton (Jonesboro)	133	5.4	27	5.1	20.3
3-4 GNR	267	10.7	78	14.7	29.2
3-5 DeKalb	319	12.8	44	8.3	13.8
4 LaGrange	160	6.4	27	5.1	16.9
5-1 South Central (Dublin)	24	1.0	<5	---	16.7
5-2 North Central (Macon)	126	5.1	27	5.1	21.4
6 East Central (Augusta)	99	4.0	20	3.8	20.2
7 West Central (Columbus)	91	3.7	26	4.9	28.6
8-1 South (Valdosta)	62	2.5	23	4.3	37.1
8-2 Southwest (Albany)	70	2.8	16	3.0	22.9
9-1 Coastal (Savannah)	148	6.0	33	6.2	22.3
9-2 Southeast (Waycross)	66	2.7	25	4.7	37.9
10 Northeast (Athens)	78	3.1	19	3.6	24.4
MSA²					
In the Atlanta MSA	1,632	65.7	312	58.8	19.1
Not in the Atlanta MSA	853	34.3	219	41.2	25.7

*Percentages may not add up to 100% due to rounding and missing data. 1. Late diagnosis: stage 3 (AIDS) diagnosis within 12 months of HIV diagnosis. Row % is percent of HIV diagnoses that were late diagnoses. 2. Atlanta Metropolitan Statistical Area (MSA) counties consist of: Barrow, Bartow, Butts, Carroll, Cherokee, Clayton, Cobb, Coweta, Dawson, DeKalb, Douglas, Fayette, Forsyth, Fulton, Gwinnett, Haralson, Heard, Henry, Jasper, Lamar, Meriwether, Morgan, Newton, Paulding, Pickens, Pike, Rockdale, Spalding, Walton.

PUBLIC HEALTH DISTRICT AND COUNTY COUNTS AND RATES

TABLE 35. HIV DIAGNOSES AND PEOPLE WITH HIV (PWH) BY PUBLIC HEALTH DISTRICT AND COUNTY (COUNTS AND RATES PER 100,000 POPULATION), GA, 2024

	HIV Diagnoses		PWH	
	N ¹	Rate/100,000 ^{2,3}	N ¹	Rate/100,000 ^{2,3}
Georgia (Total)	2,485	22.2	66,898	598.3
1-1 Northwest (Rome)	77	10.6	1,563	215.6
Bartow Co.	19	16.2	304	258.7
Catoosa Co.	<5	---	99	143.6
Chattooga Co.	<5	---	60	237.1
Dade Co.	<5	---	17	105.8
Floyd Co.	12	11.8	270	266.3
Gordon Co.	8	---	91	149.8
Haralson Co.	<5	---	45	138.9
Paulding Co.	25	13.3	508	269.4
Polk Co.	<5	---	78	174.8
Walker Co.	<5	---	91	131.2
1-2 North Georgia (Dalton)	19	3.5	872	162.8
Cherokee Co.	8	---	483	164.6
Fannin Co.	<5	---	49	189.5
Gilmer Co.	<5	---	70	210.0
Murray Co.	0	0.0	43	104.1
Pickens Co.	<5	---	48	131.2
Whitfield Co.	6	---	179	170.4
2 North (Gainesville)	64	7.9	1,242	153.2
Banks Co.	0	0.0	19	93.5
Dawson Co.	0	0.0	49	145.2
Forsyth Co.	9	---	277	98.9
Franklin Co.	<5	---	61	242.0
Habersham Co.	6	---	87	175.2
Hall Co.	28	12.6	462	208.3
Hart Co.	<5	---	56	199.6
Lumpkin Co.	7	---	46	127.7
Rabun Co.	0	0.0	20	112.9
Stephens Co.	<5	---	60	218.2
Towns Co.	0	0.0	24	182.0
Union Co.	6	---	48	173.9
White Co.	<5	---	33	111.2

TABLE 35. HIV DIAGNOSES AND PEOPLE WITH HIV (PWH) BY PUBLIC HEALTH DISTRICT AND COUNTY (COUNTS AND RATES PER 100,000 POPULATION), GA, 2024 (CONTINUED)

	HIV Diagnoses		PWH	
	N ¹	Rate/100,000 ^{2,3}	N ¹	Rate/100,000 ^{2,3}
3-1 Cobb-Douglas	195	20.8	5,015	533.8
Cobb Co.	157	19.9	4,134	524.9
Douglas Co.	38	25.0	881	580.0
3-2 Fulton	481	44.1	17,441	1599.6
Fulton Co.	481	44.1	17,441	1599.6
3-3 Clayton (Jonesboro)	133	44.7	3,437	1154.5
Clayton Co.	133	44.7	3,437	1154.5
3-4 GNR	267	21.8	5,529	451.2
Gwinnett Co.	223	22.2	4,451	443.4
Newton Co.	11	---	481	387.9
Rockdale Co.	33	33.8	597	611.6
3-5 DeKalb	319	41.4	10,302	1337.4
DeKalb Co.	319	41.4	10,302	1337.4
4 LaGrange	160	16.9	3,225	341.4
Butts Co.	26	95.8	184	678.3
Carroll Co.	17	13.1	319	245.6
Coweta Co.	15	9.5	360	227.5
Fayette Co.	15	12.0	295	235.8
Heard Co.	0	0.0	18	148.5
Henry Co.	55	21.2	1,305	503.2
Lamar Co.	0	0.0	49	236.8
Meriwether Co.	<5	---	89	415.1
Pike Co.	<5	---	18	87.1
Spalding Co.	13	18.5	285	405.5
Troup Co.	13	18.2	222	310.4
Upton Co.	<5	---	81	285.6
5-1 South Central (Dublin)	24	16.7	731	508.1
Bleckley Co.	<5	---	47	372.2
Dodge Co.	<5	---	99	504.3
Johnson Co.	<5	---	73	795.4
Laurens Co.	8	---	192	381.8
Montgomery Co.	<5	---	20	233.5
Pulaski Co.	0	0.0	51	501.0
Telfair Co.	<5	---	93	845.5
Treutlen Co.	<5	---	28	436.1
Wheeler Co.	0	0.0	35	494.6
Wilcox Co.	<5	---	93	1044.5

TABLE 35. HIV DIAGNOSES AND PEOPLE WITH HIV (PWH) BY PUBLIC HEALTH DISTRICT AND COUNTY (COUNTS AND RATES PER 100,000 POPULATION), GA, 2024 (CONTINUED)

	HIV Diagnoses		PWH	
	N ¹	Rate/100,000 ^{2,3}	N ¹	Rate/100,000 ^{2,3}
5-2 North Central (Macon)	126	22.4	2,803	497.7
Baldwin Co.	7	---	200	458.3
Bibb Co.	53	33.7	1,445	920.1
Crawford Co.	<5	---	27	218.6
Hancock Co.	<5	---	54	621.9
Houston Co.	40	22.9	582	332.8
Jasper Co.	<5	---	26	151.0
Jones Co.	<5	---	69	237.5
Monroe Co.	<5	---	81	257.7
Peach Co.	5	---	139	475.8
Putnam Co.	5	---	58	248.0
Twiggs Co.	<5	---	13	168.2
Washington Co.	<5	---	90	453.8
Wilkinson Co.	0	0.0	19	219.2
6 East Central (Augusta)	99	19.5	2,519	495.1
Burke Co.	<5	---	103	420.9
Columbia Co.	18	10.7	297	177.3
Emanuel Co.	<5	---	79	340.2
Glascock Co.	<5	---	9	---
Jefferson Co.	<5	---	61	406.2
Jenkins Co.	0	0.0	38	430.1
Lincoln Co.	0	0.0	19	236.1
McDuffie Co.	<5	---	68	312.4
Richmond Co.	65	31.5	1,740	843.4
Screven Co.	<5	---	50	349.0
Taliaferro Co.	0	0.0	10	---
Warren Co.	0	0.0	15	291.4
Wilkes Co.	<5	---	30	313.6

TABLE 35. HIV DIAGNOSES AND PEOPLE WITH HIV (PWH) BY PUBLIC HEALTH DISTRICT AND COUNTY (COUNTS AND RATES PER 100,000 POPULATION), GA, 2024 (CONTINUED)

	HIV Diagnoses		PWH	
	N ¹	Rate/100,000 ^{2,3}	N ¹	Rate/100,000 ^{2,3}
7 West Central (Columbus)	91	25.1	2,115	583.0
Chattahoochee Co.	<5	---	9	---
Clay Co.	<5	---	16	568.4
Crisp Co.	7	---	131	672.9
Dooly Co.	0	0.0	84	750.9
Harris Co.	0	0.0	51	138.1
Macon Co.	5	---	78	659.3
Marion Co.	0	0.0	15	196.5
Muscogee Co.	67	33.2	1,312	650.1
Quitman Co.	0	0.0	6	---
Randolph Co.	<5	---	42	690.2
Schley Co.	0	0.0	11	---
Stewart Co. ⁴	<5	---	140	2911.2
Sumter Co.	<5	---	151	521.2
Talbot Co.	0	0.0	30	521.3
Taylor Co.	0	0.0	36	462.5
Webster Co.	0	0.0	<5	---
8-1 South (Valdosta)	62	23.3	1,362	511.9
Ben Hill Co.	<5	---	51	301.0
Berrien Co.	<5	---	52	278.6
Brooks Co.	<5	---	81	495.6
Cook Co.	7	---	82	456.0
Echols Co.	0	0.0	16	432.7
Irwin Co.	<5	---	71	775.5
Lanier Co.	<5	---	38	363.1
Lowndes Co.	31	25.4	735	602.1
Tift Co.	10	---	181	434.0
Turner Co.	5	---	55	609.6
8-2 Southwest (Albany)	70	20.1	1,948	559.5
Baker Co.	0	0.0	7	---
Calhoun Co.	<5	---	69	1268.1
Colquitt Co.	<5	---	212	451.3
Decatur Co.	8	---	153	520.6
Dougherty Co.	35	42.5	834	1011.9
Early Co.	<5	---	32	306.4
Grady Co.	10	---	96	365.4
Lee Co.	<5	---	77	226.0
Miller Co.	0	0.0	17	291.9
Mitchell Co.	<5	---	119	585.4
Seminole Co.	0	0.0	29	312.7
Terrell Co.	0	0.0	51	586.3
Thomas Co.	<5	---	175	380.0
Worth Co.	<5	---	77	380.0

TABLE 35. HIV DIAGNOSES AND PEOPLE WITH HIV (PWH) BY PUBLIC HEALTH DISTRICT AND COUNTY (COUNTS AND RATES PER 100,000 POPULATION), GA, 2024 (CONTINUED)

	HIV Diagnoses		PWH	
	N ¹	Rate/100,000 ^{2,3}	N ¹	Rate/100,000 ^{2,3}
9-1 Coastal (Savannah)	148	21.8	3,128	461.3
Bryan Co.	9	---	63	123.3
Camden Co.	6	---	109	184.4
Chatham Co.	104	33.8	2,075	675.2
Effingham Co.	6	---	130	177.7
Glynn Co.	11	---	381	440.3
Liberty Co.	9	---	275	400.8
Long Co.	<5	---	43	210.4
McIntosh Co.	0	0.0	52	440.7
9-2 Southeast (Waycross)	66	17.2	1,491	388.2
Appling Co.	0	0.0	45	241.0
Atkinson Co.	0	0.0	20	237.1
Bacon Co.	<5	---	45	407.4
Brantley Co.	<5	---	32	170.3
Bulloch Co.	15	17.6	307	359.3
Candler Co.	<5	---	35	314.5
Charlton Co.	6	---	112	860.1
Clinch Co.	<5	---	26	378.4
Coffee Co.	10	---	158	360.3
Evans Co.	6	---	53	487.6
Jeff Davis Co.	<5	---	32	211.7
Pierce Co.	0	0.0	32	154.8
Tattall Co.	5	---	134	552.0
Toombs Co.	<5	---	99	360.2
Ware Co.	10	---	233	643.3
Wayne Co.	<5	---	128	398.4
10 Northeast (Athens)	78	13.3	1,341	228.5
Barrow Co.	17	17.7	212	220.2
Clarke Co.	29	22.3	521	400.8
Elbert Co.	<5	---	48	238.2
Greene Co.	<5	---	43	203.9
Jackson Co.	12	12.8	111	118.3
Madison Co.	<5	---	42	128.2
Morgan Co.	0	0.0	44	200.5
Oconee Co.	<5	---	26	58.1
Oglethorpe Co.	<5	---	24	148.4
Walton Co.	11	---	270	245.9

1. Counts of 1-4 are censored. 2. Rates are not calculated when the numerator is <12. 3. Persons residing in correctional facilities are included in this report and may inflate rates in certain geographic regions where there are large numbers of HIV-positive inmates. 4. Many cases associated with a federal immigration detention facility are located in this county.

DEATHS AMONG PEOPLE WITH HIV

TABLE 36. DEATHS (ALL CAUSE) AMONG PEOPLE WITH HIV (PWH), GA, 2024

	Deaths ¹ among PWH	
	N	%*
Total	874	100.0
Birth Sex		
Males	658	75.3
Females	216	24.7
Gender Identity		
Cisgender Men	646	73.9
Cisgender Women	216	24.7
Transgender Persons/Additional Gender Identity	12	1.4
Race and Ethnicity		
Black or African American, non-Hispanic	581	66.5
White, non-Hispanic	176	20.1
Hispanic or Latino, Any Race	45	5.1
Asian	0	0.0
American Indian or Alaska Native	0	0.0
Native Hawaiian or Pacific Islander	0	0.0
Multiple Races	70	8.0
Age Group at Time of Death		
<13	0	0.0
13-17	0	0.0
18-24	3	0.3
25-34	68	7.8
35-44	126	14.4
45-54	145	16.6
55-64	274	31.4
65+	258	29.5
HIV Transmission Category (Unadjusted)		
Male-to-male sexual contact (MMSC)	360	41.2
Heterosexual contact ²	159	18.2
Injection drug use (IDU)	85	9.7
MMSC and IDU	53	6.1
Adult Other ³	5	0.6
Perinatal/other pediatric ⁴	0	0.0
No risk identified or reported (NIR/NRR)	212	24.3
HIV Transmission Category (Adjusted)		
Male-to-male sexual contact (MMSC)	451	51.6
Heterosexual contact ²	252	28.8
Injection drug use (IDU)	104	11.9
MMSC and IDU	60	6.9
Adult Other ³	5	0.6
Perinatal/other pediatric ⁴	0	0.0
No risk identified or reported (NIR/NRR)	2	0.2

* Percentages may not add up to 100% due to rounding. 1. Deceased persons described in this table include those with a primary cause of death that is attributed to the underlying HIV infection or another reason (e.g. car accident, COVID-19). 2. Heterosexual contact: Defined as sexual contact with someone of the opposite sex with known risk such as injection drug use, bisexual male (applies to females only), person with hemophilia/coagulation disorder, transfusion recipient with HIV documentation, and or person with AIDS or documented HIV. 2. Includes perinatal exposure current age ≥ 13; hemophilia, blood transfusion. 3. Perinatal/other pediatric: cases <13 born to a mother who is HIV-positive, and cases <13 with other risk (e.g., sexual transmission). 4. Adjusted for missing risk using multiple imputation methods. Adjusted subtotals may be different from unadjusted subtotals due to rounding.

DISTRIBUTION OF GENERAL POPULATION BY RACE/ETHNICITY

TABLE 37. DISTRIBUTION OF THE GENERAL POPULATION BY RACE AND ETHNICITY, GA, 2024

	Population	
	N	%*
Total	11,180,878	100.0%
Race and Ethnicity		
White, non-Hispanic	5,459,978	48.8%
Black, non-Hispanic	3,580,335	32.0%
Hispanic or Latino, Any Race	1,298,399	11.6%
Asian, non-Hispanic	567,103	5.1%
American Indian or Alaskan Native	25,415	0.2%
Native Hawaiian/Pacific Islander, Non-Hispanic	8,111	0.1%
Multiracial	241,537	2.2%

* Percentages may not add up to 100% due to rounding and because not all race or ethnicity groups are represented in the table. Data source: Georgia Online Analytical Statistical Information System (<https://oasis.state.ga.us/oasis/webquery/qryPopulation.aspx>)

SUPPLEMENTARY INFORMATION

ADDITIONAL DATA ON HIV IN GEORGIA

- Georgia Department of Public Health HIV Epidemiology Unit's Main Website:
<https://dph.georgia.gov/epidemiology/hiv-epidemiology-unit/perinatal-exposure-surveillance>
 - Prior year HIV Surveillance Summary Reports are available under "HIV Case Surveillance Data Archive": <https://dph.georgia.gov/epidemiology/hiv-epidemiology-unit/hiv-case-surveillance-and-care-continuum>
 - Other resources available include data on care continuum measures, trends overtime, fact sheets, special reports, and presentations: <https://dph.georgia.gov/epidemiology/hiv-epidemiology-unit/hiv-case-surveillance-and-care-continuum>
- Centers for Disease Control and Prevention, Division of HIV Prevention:
<https://www.cdc.gov/hiv/index.html>
- AIDSvu: <https://aidsvu.org/>

REPORTING

All health care providers diagnosing and/or providing care to a patient with HIV are required by Georgia law (O.C.G.A. §31-12-1) to report HIV infection using the HIV/AIDS Case Report Form.

Case report forms should be completed within seven (7) days of diagnosing a patient with HIV and/or AIDS or within seven (7) days of assuming care of an HIV positive patient who is new to the provider, regardless of whether the patient has previously received care elsewhere.

Adult and Pediatric case report forms are available at <https://dph.georgia.gov/epidemiology/hiv-epidemiology-unit/hiv-and-aids-case-reporting>

For more questions on HIV case reporting in Georgia please contact the HIV Surveillance Coordinator at 1-800-827-9769 (no faxing permitted). eFax: 404-506-9297

REQUESTING DATA

To request data, please visit <http://dph.georgia.gov/phip-data-request> to create a PHIP data request account and login.

