

03/31/24

Demographic and Clinical Characteristics of Selected Subgroups of Persons with Diagnosed HIV Infection

Georgia Medical Monitoring Project, 2015–2021



The **Georgia Medical Monitoring Project: Demographic and Clinical Characteristics of Selected Subgroups of Persons with Diagnosed HIV Infection, 2015–2021** is published by the Georgia Department of Public Health (DPH), HIV Epidemiology Section, 200 Piedmont Ave, SE / West Tower, Suite 1002, Atlanta, Georgia 30334.

The Georgia Medical Monitoring Project: Demographic and Clinical Characteristics of Selected Subgroups of Persons with Diagnosed HIV Infection 2015–2021 is not copyrighted and may be used and reproduced without permission. Citation of the source is, however, appreciated.

SUGGESTED CITATION: Georgia Department of Public Health, HIV Epidemiology Section *Medical Monitoring Project: Demographic and Clinical Characteristics of Selected Subgroups of Persons with Diagnosed HIV Infection, Georgia Medical Monitoring Project, 2015–2021*, <https://dph.georgia.gov/georgia-medical-monitoring-project-mmp>, Published 3/31/2024, [Accessed: date]

ACKNOWLEDGEMENTS: Publication of this report was made possible with the contributions of the Georgia Medical Monitoring Project (MMP) Team, the national MMP staff at the Centers for Disease Control and Prevention (CDC), assistance from HIV care facilities, and the cooperation of participants.

Georgia MMP Team: Natalie Lucas, MPH; Kiswana Branch, MPH; Bola Griffin, MPH; Melissa Gousse, MPH; Timothy Lockhart, MPH; Shelbie Richardson MPH; and Stephen Ray, MSPH.

The report was prepared using code provided by CDC to replicate the national report, and the text is based on the national report: *Centers for Disease Control and Prevention. Behavioral and Clinical Characteristics of Persons with Diagnosed HIV Infection—Medical Monitoring Project, United States, 2021 Cycle (June 2021–May 2022). HIV Surveillance Special Report 32*. <https://www.cdc.gov/hiv/library/reports/hiv-surveillance.html>. Published May 2023. Accessed 08/31/2023.

This report was prepared by the following staff of the Georgia Department of Public Health: Stephen Ray, MSPH; Jenna Gettings, DVM, MPH; Cherie Drenzek, DVM, MS.

TABLE OF CONTENTS

COMMENTARY.....	4
TABLE 1. SELECTED DEMOGRAPHICS OF PEOPLE WITH HIV —MEDICAL MONITORING PROJECT, GEORGIA, 2015–2021	5
FIGURE 1: SEXUAL PREFERENCE, GEORGIA MMP 2015–2021	6
FIGURE 2: EDUCATION LEVEL, GEORGIA MMP 2015–2021	7
FIGURE 3: ANY DISABILITY, GEORGIA MMP 2015–2021	7
FIGURE 4: CURRENTLY EMPLOYED, GEORGIA MMP 2015–2021	8
FIGURE 5: YEARLY INCOME, GEORGIA MMP 2015–2021	8
TABLE 2. HEALTH PROVIDER AND PUBLIC ASSISTANCE PROGRAM USAGE BY PEOPLE WITH HIV —MEDICAL MONITORING PROJECT, GEORGIA, 2015–2021	9
FIGURE 6: PRIVATE HEALTH INSURANCE, GEORGIA MMP 2015–2021.....	10
FIGURE 7: RECEIVED CARE AT A RYAN WHITE FACILITY IN THE PAST 12 MONTHS, GEORGIA MMP 2015–2021.....	11
TABLE 3. HOSPITAL AND EMERGENCY DEPARTMENT USAGE BY PEOPLE WITH HIV IN THE PAST TWELVE MONTHS, GEORGIA, 2015–2021	12
FIGURE 8: NUMBER OF EMERGENCY DEPARTMENT VISITS, GEORGIA MMP 2015–2021	13
TABLE 4. SEXUALLY TRANSMITTED INFECTION (STI) SCREENING IN THE PAST TWELVE MONTHS AMONG PEOPLE WITH HIV, GEORGIA, 2015–2021.....	14
FIGURE 9: SCREENED FOR GONORRHEA AND CHLAMYDIA, GEORGIA MMP 2015–2021.....	15
FIGURE 10: SCREENED FOR SYPHILIS, GEORGIA MMP 2015–2021	15
FIGURE 11: SCREENED FOR ALL THREE STI, GEORGIA MMP 2015–2021	16
TABLE 5. HIV STATUS, SUPPRESSION, AND CARE RETENTION AMONG PEOPLE WITH HIV, GEORGIA, 2015–2021	17
TABLE 6. ART ADHERENCE AMONG PEOPLE WITH HIV, GEORGIA, 2015–2021.....	19
TABLE 7. REASONS FOR MISSING LAST ART DOSE AMONG PEOPLE WITH HIV WHO INDICATED MISSING LAST ART DOSE IN THE PAST THIRTY DAYS, GEORGIA, 2015–2021	21
FIGURE 12: NUMBER OF REASONS GIVEN FOR MISSING LAST ART DOSE IN THE PAST THIRTY DAYS, GEORGIA MMP 2015–2021	22



COMMENTARY

The purpose of this report is to describe selected demographic and clinical characteristics of Black males, White males, and Black females, three groups surveyed that account for 86% of the Medical Monitoring Project (MMP) participants and approximately 81% of persons with HIV in Georgia. Findings are also stratified by location both for residents of the Atlanta Metropolitan Statistical Area (MSA) and for persons residing in other parts of the state (referred to as Atlanta and non-Atlanta residents). This report uses MMP data from the 2015 through 2021 cycles (i.e., June 2015–May 2021).

The Georgia MMP is a Centers for Disease Control and Prevention (CDC)-funded surveillance project aimed at assessing the clinical and behavioral characteristics of people with HIV (PWH) in Georgia. Every year, 500 PLWH are randomly sampled from the Georgia HIV Surveillance Registry and are assigned to the MMP team at the Georgia Department of Public Health (GDPH) to be contacted to determine eligibility and interest in being interviewed. In addition, a medical record abstraction is performed with the participant's consent in order to gather the clinical data as part of the project. MMP methods are described in greater detail in the 2015–2021 Georgia MMP Surveillance Report which can be found at: <https://dph.georgia.gov/epidemiology/georgias-hiv-aids-epidemiology-section/ga-mmp>.

Technical Notes:

Of the 3,500 total PWH sampled during the 2015 to 2021 data cycles, 1,369 (39%) agreed to an interview and medical record abstraction. Data are weighted to take into account the sampling scheme; all percentages presented are weighted percentages. Most of the information shown in the tables, including gender and race/ethnicity, is from the interview, but the following pieces of information are from the chart abstractions (either in tandem with interview information or solely): viral suppression status, retention in HIV care, CD4 counts, and sexually transmitted infection (STI) testing. Participants are categorized as Atlanta MSA or non-Atlanta MSA residents based on the location of the clinic where they primarily received HIV care at the time of interview. Location of care was available for approximately 85% of participants in this analysis. Because MMP is drawn from the HIV surveillance registry, it is not restricted to persons who are in care. Persons who are not in care are, however, hard to reach and may be underrepresented among respondents. Thus, estimates of retention in care and viral suppression may be overestimated.

All figures that present notable differences between Atlanta residents and non-Atlanta residents are restricted to Black men and women due to the relatively small number of White male PWH residing outside of the Atlanta MSA.

Table 1. Selected Demographics of People with HIV - Medical Monitoring Project, Georgia, 2015–2021


	Black Males n=698		White Males n=191		Black Females n=295	
	wt'd %	n	wt'd %	n	wt'd %	n
Sexual preference						
Lesbian or gay	51.5	349	74.3	138	1.1	4*
Heterosexual or straight	31.2	224	13.3	26	94.5	275
Bisexual	13.9	92	11.9	23	3.3	9*
Other sexual orientation	3.5	24	0.5	1*	1.0	3*
Age at interview						
18–29 years	14.8	98	3.6	8*	8.2	23
30–39 years	24.2	167	10.6	18	18.7	46
40–49 years	20.8	141	15.9	32	26.9	77
≥50 years	40.1	292	69.8	133	46.3	149
Time since HIV diagnosis						
<5 years	21.2	148	12.5	22	15.1	43
5–9 years	23.9	165	12.4	22	26.5	74
≥10 years	54.8	385	75.1	147	58.4	178
Education level						
Less than high school	10.6	77	6.3	11*	24.6	78
High school diploma or GED	26.5	177	13.4	27	30.9	88
More than high school	62.9	439	80.3	151	44.5	126
Any disability						
Yes, has any disability	31.4	220	38.5	69	53.2	158
Currently employed						
Yes, currently employed	57.1	393	67.9	128	43.5	127
Yearly income						
\$0–\$19,999	44.0	274	27.1	49	62.9	169
\$20,000–\$39,999	25.6	165	23.9	41	21.6	59
\$40,000–\$74,999	22.0	141	27.4	50	11.1	25
≥ \$75,000	8.4	58	21.6	40	4.5	10*

*: A count with an asterisk indicates a corresponding percentage with a coefficient of variation ≥ 0.30 and should be interpreted with caution.

Counts may not add up to the reported number of participants (n) due to missing information

All percentages shown are weighted percentages

FIGURE 1: SEXUAL PREFERENCE, GEORGIA MMP 2015–2021



Overall, approximately half of Black men and three quarters of White men reported being gay, while almost all Black women reported being heterosexual (Table 1). Among Black men, the majority of Atlanta residents identified as gay while the majority of non-Atlanta residents identified as heterosexual (Figure 1). Due to few people with HIV identifying as such, approximate percentages shown for “Other sexual orientation” across all groups and all orientations except “Heterosexual, or straight” for Black females should be interpreted with caution.


Age at time of interview:

Across all groups, there were a higher number of older people with HIV than younger people with HIV (Table 1). The largest sub-group across all groups was the ≥50 years sub-group. There were no noteworthy large differences comparing Black Atlanta residents and non-Atlanta residents for this measurement (These data are not presented in figures).

Time since HIV diagnosis:


Overall, approximately half of Black men and three quarters of White men were diagnosed more than ten years ago (Table 1). 58% of Black women were diagnosed more than ten years ago. There were no noteworthy large differences comparing Black Atlanta residents and non-Atlanta residents for this measurement (These data are not presented in figures).

FIGURE 2: EDUCATION LEVEL, GEORGIA MMP 2015–2021




Overall, approximately 60% of Black males and 80% of White males had pursued further education past a high school diploma or GED (Table 1). Among Black females, only about 45% of Black females had pursued further education. Educational attainment was lower among non-Atlanta residents than Atlanta residents, regardless of race or gender (Figure 2). Of interest, about 7% of Black males living in Atlanta had less than a high school education, while 28% of Black males living outside of the Atlanta area had less than a high school education.

FIGURE 3: ANY DISABILITY, GEORGIA MMP 2015–2021




Overall, about a third of Black males and 39% of White males self-reported that they had a disability. Over half of Black females self-reported a disability (Table 1). A higher percentage of Black male non-Atlanta residents self-reported disabilities compared to Black male Atlanta residents (Figure 3). This same magnitude of difference was not observed when comparing between Black female Atlanta residents and non-residents.

FIGURE 4: CURRENTLY EMPLOYED, GEORGIA MMP 2015–2021



Overall, approximately 57% of Black males and 68% White males reported that they were currently employed. Only approximately 44% of Black females reported that they were currently employed (Table 1). Among Black males, current employment was higher among Atlanta residents than non-Atlanta residents. Similarly, current employment was higher among Black female Atlanta residents than non-Atlanta residents. (Figure 4).

FIGURE 5: YEARLY INCOME, GEORGIA MMP 2015–2021



Across both groups of Black male and females, the largest sub-group was those earning between \$0–\$19,999. The majority of Black females earned below \$20,000 (Table 1). Yearly income was lower on average for non-Atlanta residents compared to Atlanta residents, especially among Black females (Figure 5). Approximate percentages shown for Black women and Black male non-Atlanta residents for the two highest income levels (\$40,000–\$74,999 and ≥\$75,000) should be interpreted with caution due to low response rate.

Table 2. Health Provider and Public Assistance Program Use by People with HIV —Medical Monitoring Project, Georgia, 2015–2021

	Black Males n=698		White Males n=191		Black Females n=295	
	wt'd %	n	wt'd %	n	wt'd %	n
Type of health insurance or coverage¹ for antiretroviral therapy, past 12 months						
Ryan White/ADAP ²	45.7	330	34.8	69	38.9	117
Medicaid	20.1	169	14.5	24	56.5	159
Private health insurance	36.9	283	52.8	103	26.4	71
Medicare	22.9	162	39.3	70	30.8	96
Care at a Ryan White facility in the past 12 months						
Yes	72.8	482	54.3	101	78.0	225
Received SSI³ in the past 12 months						
Yes, received SSI	16.4	118	13.8	23	32.0	93
Received SSDI⁴ in the past 12 months						
Yes, received SSDI	20.2	141	26.2	46	28.3	84
Went without food in the past 12 months						
Yes, went without food	20.8	137	11.7	22	14.4	46

Counts may not add up to the reported number of participants (n) due to missing information

All percentages shown are weighted percentages

1: Types of health insurance are not mutually exclusive, and people with HIV may report multiple types of health insurance or coverage

2: AIDS Drug Assistance Program

3: SSI: Supplemental Security Income


4: SSDI: Social Security Disability Insurance

Type of health insurance or coverage for antiretroviral therapy (ART), past 12 months

A third of all White males compared to 45% of Black males reported utilizing Ryan White/ADAP coverage for their ART and HIV care. For Black females, approximately 40% reported utilizing Ryan White/ADAP coverage. About 57% of Black females with HIV reported having Medicaid coverage, compared to 20% of Black males and approximately 15% of White males. About 40% of White males reported being covered under Medicare compared to 23% of Black males and 31% of Black females.


There were no noteworthy large differences comparing Black Atlanta residents and non-Atlanta residents for this measurement (These data are not presented in figures).

FIGURE 6: PRIVATE HEALTH INSURANCE, GEORGIA MMP 2015–2021



Approximately 40% of Black males and half of White males carried some form of private insurance. Only about a quarter of Black females carried private insurance (Table 2). Non-Atlanta Black male residents were less likely to have private insurance compared to Atlanta Black male residents. About 39% of Atlanta Black female residents carried private insurance while only 17% of non-Atlanta Black female residents carried private insurance.

FIGURE 7: RECEIVED CARE AT A RYAN WHITE FACILITY IN THE PAST 12 MONTHS, GEORGIA MMP 2015–2021



A higher proportion of Black males and females received care at Ryan White facilities in the past 12 months compared with White males (Table 2). A higher proportion of Black non-Atlanta residents compared to Black Atlanta residents received care at Ryan White clinics, regardless of gender (Figure 7).

Received Supplemental Security Income (SSI) in the past 12 months:

Most people with HIV, regardless of race or gender, did not receive SSI (Table 2). A higher proportion of Black females received SSI compared to Black and White males, but overall usage remained low (32%). There were no noteworthy large differences comparing Black Atlanta residents and non-Atlanta residents for this measurement (These data are not presented in figures).

Received Social Security Disability Insurance (SSDI) in the past 12 months:

About 20% of Black males received SSDI benefits in the past 12 months; about a quarter of White males and Black females received SSDI benefits in the past 12 months (Table 2). There were no noteworthy large differences comparing Black Atlanta residents and non-Atlanta residents for this measurement (These data are not presented in figures).

Went without food in the past 12 months:


Most people with HIV, regardless of race or gender, reported that they did not go without food in the past 12 months (Table 2). A higher proportion of Black males (20%) went without food compared to Black females (14%) and White males (12%), but the proportion is low. There were no noteworthy large differences comparing Black Atlanta residents and non-Atlanta residents for this measurement (These data are not presented in figures).

Table 3. Hospital and Emergency Department Use by People with HIV in the Past Twelve Months, Georgia, 2015–2021

	Black Males n=698		White Males n=191		Black Females n=295	
	<u>wt'd %</u>	<u>n</u>	<u>wt'd %</u>	<u>n</u>	<u>wt'd %</u>	<u>n</u>
Number of visits to the emergency department						
0	58.2	402	58.9	117	51.9	148
1	19.2	133	28.7	49	15.8	47
≥2	22.6	157	12.4	23	32.3	92
Number of hospital admissions						
0	80.7	559	80.0	152	75.8	215
1	12.5	85	16.2	29	16.2	49
≥2	6.7	48	3.8	8*	8.1	23

*: A count with an asterisk indicates a corresponding percentage with a coefficient of variation ≥ 0.30 and should be interpreted with caution.
 Counts may not add up to the reported number of participants (n) due to missing information
 All percentages shown are weighted percentages

FIGURE 8: NUMBER OF EMERGENCY DEPARTMENT VISITS,
GEORGIA MMP 2015–2021



Around 45% of people used emergency room services. Approximately half of Black females visited the Emergency Department (ED) at least once compared to about 40% of Black males or White males (Table 3). The proportion of Black people with HIV living outside of Atlanta who visited the ER two or more times was higher compared with Black people with HIV that were Atlanta residents (Figure 8).

Number of Hospital Admissions:

Approximately 20% of Black males and White males, and a quarter of Black females had one or more hospital admissions during the previous 12 months.


There were no noteworthy large differences comparing Black Atlanta residents and non-Atlanta residents for this measurement (These data are not presented in figures).

Table 4. Sexually Transmitted Infection (STI) Screening in the Past Twelve Months among People with HIV, Georgia, 2015–2021

	Black Males n=698		White Males n=191		Black Females n=295	
	wt'd %	n	wt'd %	n	wt'd %	n
Screened for gonorrhea						
Yes, screened	52.6	338	39.2	67	46.8	130
Screened for chlamydia						
Yes, screened	52.6	338	39.2	67	46.8	130
Screened for syphilis						
Yes, screened	67.7	443	65.8	114	58.9	164
Screened for all three STIs						
Yes, screened	47.4	303	38.4	66	39.3	107

Counts may not add up to the reported number of participants (n) due to missing information
All percentages shown are weighted percentages


FIGURE 9: SCREENED FOR GONORRHEA AND CHLAMYDIA, GEORGIA MMP 2015–2021



Only about 39% of White males were screened for gonorrhea and chlamydia compared to 52% of Black males and 47% of Black females (Table 4). The proportion tested was only slightly higher (1-4 percentage points) when restricting to those reporting sexual activity (not shown). Screening rates for gonorrhea and chlamydia were lower across both Black males and Black females for non-Atlanta residents compared to Atlanta residents (Figure 9).


Note: This difference seen in screening rates are likely due to age distribution across each subgroup.

FIGURE 10: SCREENED FOR SYPHILIS, GEORGIA MMP 2015–2021



Screening for syphilis was higher than testing for gonorrhea and chlamydia: about 65% of Black males and White males were tested for syphilis and about 60% of Black females were tested (Table 4). There was no difference when restricting to those reporting sexual activity (not shown). Screening for syphilis was lower among Black males outside of Atlanta compared to Atlanta residents (Figure 10).

FIGURE 11: SCREENED FOR ALL THREE STI, GEORGIA MMP
2015–2021



Screening for all three STIs follows the same pattern observed in screening for gonorrhea and chlamydia. There was a lower percentage of non-Atlanta residents being tested for all three STIs compared to Atlanta residents (Figure 11), and there was no noteworthy difference in proportion among those reporting sexual activity (not shown).

Table 5. HIV Status, Suppression, and Care Retention among People with HIV, Georgia, 2015–2021

	Black Males n=698		White Males n=191		Black Females n=295	
	wt'd %	n	wt'd %	n	wt'd %	n
HIV infection- Stage 3 (AIDS)¹						
AIDS (Stage 3 infection)	52.4	385	50.0	101	59.2	184
Ever taken ART						
Yes, taken ART	97.2	676	98.6	187	98.3	287
Currently taking ART						
Yes, taking ART	91.0	638	92.9	178	93.2	273
Lowest CD4 count in the past 12 months						
<200 cells/μL	12.4	75	6.1	11	13.4	33
200–499 cells/μL	34.2	196	38.4	61	31.6	82
≥500 cells/μL	53.4	301	55.5	78	55.0	137
Viral suppression²						
Most recent viral load test result <200/mL	67.7	479	77.0	146	68.9	209
Durable viral suppression³						
All viral load test results <200 copies/mL	58.3	418	74.3	140	60.7	184
Retained in care⁴, past 12 months						
Yes	78.8	538	79.3	147	81.7	242
Retained in care, past 24 months						
Yes	57.2	386	65.1	120	65.0	189

1: HIV infection- Stage 3 at time of interview


2: Viral suppression defined as HIV viral load of less than 200 count/mL of blood, using nucleic acid testing (NAT)

3: Durable viral suppression defined as all viral load tests performed being less than 200 count/mL during the past 12 months before the interview. Quantity of tests are not considered.

4: Retention in care was defined as at least two elements of outpatient HIV care at least 90 days apart in each 12-month period. Outpatient HIV care was defined as any documentation in the medical record of the following: encounter with an HIV care provider, viral load test result, CD4 test result, HIV resistance test or tropism assay, ART prescription, PCP prophylaxis, or MAC prophylaxis.

Counts may not add up to the reported number of participants (n) due to missing information

All percentages shown are weighted percentages



HIV Infection Stage 3 (AIDS):

Approximately half of Black males and White males had Stage 3 HIV (AIDS) at the time of interview. For Black females, about 60% had Stage 3 HIV (Table 5).

There were no noteworthy large differences comparing Black Atlanta residents and non-Atlanta residents for this measurement (These data are not presented in figures).

Ever taken and currently taking ART:

Almost all people with HIV had ever taken some form of ART (Table 5), and the vast majority were on ART at the time of their interview.

There were no noteworthy large differences comparing Black Atlanta residents and non-Atlanta residents for this measurement (These data are not presented in figures).

Lowest CD4 count in the past 12 months:

Overall, about 50% of people with HIV across all groups had CD4 counts ≥ 500 cells/ μ L, and less than 14% had a CD4 count of < 200 cells/ μ L (Table 5). About 6% of White males had a CD4 count of < 200 cells/ μ L, while approximately 13% of Black males and females both had a CD4 count of < 200 cells/ μ L.

The proportion that had counts of < 200 cells/ μ L was not different for Black Atlanta and non-Atlanta residents (These data are not presented in figures).

Viral suppression:

Viral suppression was achieved by approximately two-thirds of Black males and females and 80% of White males in their most recent viral load test (Table 5).

There were no noteworthy large differences comparing Black Atlanta residents and non-Atlanta residents for this measurement (These data are not presented in figures).

Durable viral suppression for the past 12 months:

The proportion of persons with HIV achieving durable viral suppression (all viral load tests < 200 copies/mL in the past 12 months) was lower than those achieving recent viral suppression. Around 60% of Black females and males had viral loads of < 200 copies/mL at every test in the past 12 months compared with three-quarters of White males (Table 5).

There were no noteworthy large differences comparing Black Atlanta residents and non-Atlanta residents for this measurement (These data are not presented in figures).

Retained in Care for the Past 12/24 Months:

Retention in care was defined as at least two elements of outpatient HIV care at least 90 days apart in each 12-month period. Outpatient HIV care was defined as any documentation in the medical record of the following: encounter with an HIV care provider, viral load test result, CD4 test result, HIV resistance test or tropism assay, ART prescription, PCP prophylaxis, or MAC prophylaxis.

For the past 12 months, most people with HIV were retained in care. About 80% across all three groups observed were retained in care (Table 5).


When expanding the observation period to the past 24 months, only approximately 57% of Black males and 65% of White males and Black females were retained in care (Table 5).

There were no noteworthy large differences comparing Black Atlanta residents and non-Atlanta residents for both measurements (both past 12 months and past 24 months) (These data are not presented in figures).

Table 6. ART Adherence among People with HIV, Georgia, 2015–2021

	Black Males n=698		White Males n=191		Black Females n=295	
	<u>wt'd %</u>	<u>n</u>	<u>wt'd %</u>	<u>n</u>	<u>wt'd %</u>	<u>n</u>
<u>In the past 30 days:</u>						
<i>Question: How many days did you miss at least 1 dose of any of your HIV medicine(s)?</i>						
0	53.9	344	70.7	125	60.7	166
≥1	46.1	293	29.3	53	39.3	106
<i>Question: How well did you do at taking your HIV medicine(s) in the way you were supposed to?</i>						
Very poor	1.0	7*	0.0	0	1.2	3*
Poor	2.5	15	0.0	0	1.9	6*
Fair	5.7	33	1.2	3*	5.3	15
Good	12.6	82	11.1	17	13.9	39
Very good	28.7	185	26.2	47	26.7	73
Excellent	49.5	316	61.4	111	51.0	137
<i>Question: How often did you take your HIV medicine(s) in the way you were supposed to?</i>						
Never	0.6	4*	0.0	0	0.7	2*
Rarely	0.5	3*	0.0	0	2.6	6*
Sometimes	2.4	17	0.0	0	1.8	7*
Usually	4.9	31	4.6	6*	3.3	10*
Almost always	28.9	182	27.5	49	27.1	69
Always	62.7	401	67.9	123	64.4	179
<i>Question: How often were you troubled by ART side effects?</i>						
Never	73.4	467	70.1	126	68.6	189
Rarely	13.1	88	17.0	28	18.7	44
About half the time	6.7	39	7.1	12	5.6	16
Most of the time	3.5	23	2.3	5*	4.9	13
Always	3.4	19	3.5	6*	3.1	8*

Counts may not add up to the reported number of participants (n) due to missing information
All percentages shown are weighted percentages



ART Adherence in the Past 30 Days:

About 39% of Black females and 29% of White males reported missing doses of their HIV medication on at least one day in the past 30 days while about 46% of Black males reported missing at least one dose (Table 6). There were no noteworthy large differences comparing Black Atlanta residents and non-Atlanta residents for this measurement (These data are not presented in figures).

When asked how well they take their HIV medication as they are supposed to, about three-quarters of Black males and females and approximately 87% of White males reported "Very Good" or "Excellent". When asked how often they took their HIV medication correctly, about 90% of people with HIV reported "Almost Always" or "Always" (Table 6).

There were no noteworthy large differences comparing Black Atlanta residents and non-Atlanta residents for this measurement (These data are not presented in figures).

Overall, about 85% of Black males and females reported "Never" or "Rarely" being troubled by side effects and 90% of White males reported "Never" or "Rarely" being troubled by side effects. About 13% of people with HIV reported having side effects more than rarely. (Table 6).

There were no noteworthy large differences comparing Black Atlanta residents and non-Atlanta residents for this measurement (These data are not presented in figures).

Table 7. Reasons for Missing Last ART Dose among People with HIV who Indicated Missing Last ART Dose in the Past Thirty Days, Georgia, 2015–2021

	Black Males n=293		White Males n=53		Black Females n=106	
	wt'd %	n	wt'd %	n	wt'd %	n
Reason for last missed dose:						
<i>Change in your daily routine or were out of town</i>						
Yes	47.5	145	47.6	25	35.6	37
<i>Fell asleep or overslept</i>						
Yes	39.5	117	46.4	24	46.8	50
<i>Forgot to take HIV medicines</i>						
Yes	66.6	196	64.6	33	57.6	59
<i>Had a problem getting a prescription, a refill, insurance coverage, or paying for HIV medicine(s)</i>						
Yes	27.9	80	10.3	5*	18.2	21
<i>Felt depressed or overwhelmed</i>						
Yes	15.7	45	12.6	5*	15.7	19
<i>Had side effects from your HIV medicine(s)</i>						
Yes	9.8	27	9.1	5*	18.5	17
<i>Number of reasons¹ given for missing last ART dose</i>						
0 reasons ²	2.8	9*	4.5	2*	4.4	6*
1 reason	35.0	103	31.2	19	42.7	45
2-3 reasons	51.1	145	58.0	29	40.3	42
4-6 reasons	11.1	36	6.4	3*	12.5	13

*: A count with an asterisk indicates a corresponding percentage with a coefficient of variation ≥ 0.30 and should be interpreted with caution.

1: Six responses were chosen but do not cover all possible responses to reasons why missed ART. Several reasons were skipped due to overall low response rates

2: This response is possible if participant does not give a confirmatory response to any of the six highlighted questions; another reason can be one of the reasons why ART was missed was not one of the six reasons highlighted in this table.

Counts may not add up to the reported number of participants (n) due to missing information

All percentages shown are weighted percentages




Reason(s) for missing last ART dose among those who missed their last dose, past 30 days:

The most common reasons for missing their last ART dose included changes in daily routine, falling asleep too early or oversleeping, and forgetting to take their medication. About 10% of Black males and White males reported issues with side effects related to their ART medication, while 18% of Black females reported side effects. About 15% reported feeling depressed or overwhelmed. A higher proportion of Black males (28%) and females (18%) than White males (10%) reported issues with ART medication prescriptions (Table 7).

Either there were no noteworthy large differences comparing Black Atlanta residents and non-Atlanta residents for this measurement or the number of confirmatory responses was too small given this selection of participants to accurately describe the population by Atlanta metropolitan status (These data are not presented in figures).

FIGURE 12: NUMBER OF REASONS GIVEN FOR MISSING LAST ART DOSE IN THE PAST THIRTY DAYS, GEORGIA MMP 2015–2021



An overwhelming majority of people with HIV who had missed doses of HIV medication on at least one day answered yes to at least one reason for missing their ART dose. About 43% of Black females, 31% of White males, and a third of Black males reported only one reason why they missed their last ART dose. The majority of males responded having 2-3 reasons why they missed their last ART dose, with 51% of Black males and 58% of White males responding as such. 40% of Black females reported 2-3 reasons (Table 7). Overall, a higher proportion of White males responded having 2-3 reasons for missing an ART dose, while a higher proportion of Black females responded having either one reason or ≥4 reasons given for missing an ART dose.

Either there were no noteworthy large differences comparing Black Atlanta residents and non-Atlanta residents for this measurement or the number of confirmatory responses was too small given this selection of participants to accurately describe the population by Atlanta metropolitan status (These data are not presented in figures).