

**Report on Stroke in Georgia 2019, as Required by the
Coverdell-Murphy Act, Georgia SB549, Amended by Georgia HB853
Compiled by the Georgia Coverdell Acute Stroke Registry
Georgia Department of Public Health
December 2019**

Background

Why should we care about stroke in Georgia?

- **Georgia's age-standardized stroke death rate in 2018 was 14.5 percent higher** than the national average.¹
- **In 2018, Georgia had the seventh highest stroke death rate** in the U.S.¹
- **Stroke is the fourth leading cause of death** in Georgia (**4,553 stroke deaths in 2018**).¹
- **In 2018, about 17 percent of Georgia stroke deaths were premature** (i.e., among persons under the age of 65 years).¹
- **In 2018, the age-adjusted stroke death rate for Blacks in Georgia was 53.7 per 100,000 population, which was 33 percent higher** than the rates for Whites.¹
- Stroke is a **leading cause of disability**.² Treatment of eligible stroke patients with the drug Alteplase (a tissue plasminogen activator) can reduce disability by 30 percent, but the drug needs to be administered in the first three hours after symptom onset.³
- **In 2018, Georgians had more than 22,000 stroke hospitalizations**
 - The median charge per hospitalization was around **\$37,570**.
 - The total stroke-related hospitalization charges **were over \$1.5 billion in Georgia**.
- **Georgia is in the "Stroke Belt,"** an area in the southeastern U.S. with stroke death rates that are approximately **30 percent higher** than the rest of the U.S. The coastal plains of Georgia are in the "buckle" of the Stroke Belt, an area with stroke death rates about **40 percent higher** than the rest of the nation.⁴
 - The higher death rates seen in the Stroke Belt can be collectively explained, in large part, by demographic and socioeconomic factors and the prevalence of stroke risk factors and chronic diseases like diabetes and hypertension.⁵

- **In 2018, only 63 percent** of adult Georgians knew all three signs of stroke – facial droop, arm weakness, and slurred speech – and the importance of calling 911 immediately.
- **Georgia stroke patients have higher prevalence rates for stroke-related risk factors versus adult Georgians.** The 2017/18 Behavioral Risk Factor Surveillance System (BRFSS) and 2018 GCASR data showed:⁶

Risk Factor	Acute Stroke Patients (%) ^a	Adult Georgians (%)
Hypertension	79.7	33.1
Dyslipidemia	43.8	31.1
Diabetes Mellitus	36.2	12.6
Coronary Artery Disease/Prior Myocardial Infarction	21.8	7.1
Atrial Fibrillation/Flutter	14.4	--
Smoking	21.4	16.1
Physical Inactivity [§]	--	31.0

a: GCASR 2018

§: No physical activity or exercise during the past 30 days other than their regular job.

Coverdell-Murphy Act Required Reporting

To assure that patients are receiving the appropriate level of care and treatment at each level of stroke center, Georgia's Coverdell-Murphy Act (CMA), Senate Bill 549, enacted in 2008 and amended in 2016 (House Bill 853), requires the reporting of stroke care related data to the Georgia Department of Public Health (DPH) as part of the Georgia Coverdell Acute Stroke Registry (GCASR).^{7,8} The required data elements are used to generate performance measures to monitor the quality of stroke care among the designated stroke centers. GCASR currently has 80 participating acute care hospitals, of which 4 are designated as comprehensive, 43 as primary, and 19 as remote treatment stroke centers (Map 1). In 2018, the designated hospitals received 89 percent of Georgians admitted with acute stroke across the state.

Summary of Data Findings

Based on 2008-2017 hospital discharge data from 12 designated Remote Treatment Stroke Centers, acute ischemic stroke patients **treated after the hospitals were designated had 47 percent lower odds of in-hospital death** compared to patients **admitted when the hospitals were not participating** in the Georgia Coverdell Acute Stroke Registry, which aims to improve the quality of stroke care.

From 2016 to 2018, Georgians received high quality stroke care consistently.

- **Nine out of ten** eligible ischemic stroke patients received the clot-busting drug Alteplase intravenously.
- The median time from hospital arrival to the administration of Alteplase was **shortened from 52 minutes** in 2016 **to 50 minutes** in 2018.
- About **43 percent** of the patients received Alteplase in the first 45 minutes of hospital arrival in 2018 **compared to 39 percent** in 2016.

Numbers for other quality indicators, such as venous thromboembolism prophylaxis, antithrombotic medication, stroke education and discharge on appropriate medication, remained **consistently high (above 90 percent)** from 2016 to 2018.

However, **less than 60 percent** of stroke patients were transported to hospitals by EMS and the proportion of patients who arrived at the hospital in the first two hours of symptom onset remained persistently **below 40 percent**, indicating the need to raise public awareness to identify stroke in the community, call 911, and transport patients to designated centers rapidly.

Discussion

The Centers for Disease Control and Prevention (CDC) funds the Georgia Coverdell Acute Stroke Registry (GCASR) to improve stroke systems of care in Georgia. Participating EMS agencies, hospitals, and post-acute care facilities are working to strengthen the existing relationships and developing new approaches to deliver the best stroke care possible at all levels of the patient care continuum. Currently, 45 EMS agencies and 80 acute care facilities in Georgia, of which 19 hospitals were designated as Remote Treatment Stroke Centers, participate in the GCASR, and they have already had a major impact on the lives of Georgians by reducing mortality and limiting disability from stroke.^{9,10}

Shortening the time between symptom onset and hospital arrival is crucial for better patient outcomes. Currently, 64 percent of patients arrived at hospitals two hours or later after symptom onset and 43 percent transport themselves to a hospital; therefore, it is critical that all stakeholders exert a concerted effort to increase public awareness about the signs of acute stroke and the importance of calling 911 for a swift transfer of patients to one of the designated stroke centers for the appropriate level of treatment.

Moreover, we must continue to improve stroke prevention and treatment across the state by reducing the prevalence of stroke risk factors in Georgia. Adults 55 years and older have a higher risk for stroke.¹¹ Based on the National Center for Health Statistics projection, 16 percent of Georgia residents are expected to be 65 years and older by the 2030.¹² Thus, the number of Georgians affected by stroke is expected to rise over the next decade. This will increase costs, both financially and in terms of productive years of life lost.

Stroke is a major cause of disability, and adequate post-hospital care contributes significantly to reducing late complications of the acute incident. It is imperative, therefore, to monitor the quality of post-hospital stroke care. With the current five-year grant from the CDC, the GCASR has expanded its reach and the registry adopted performance indicators to improve the quality of post-hospital.

Table 1. Performance Measures for Designated Stroke Centers, GCASR, 2015-2017

Performance Measure	2016	2017	2018
Reach			
% of state acute stroke admissions in designated GCASR hospitals	87	87	89
Public Awareness			
% of patients presenting to ED with acute stroke or TIA that arrive by EMS	57	57	57
% of patients with acute stroke or TIA who arrive at ED in less than 2 hours from time last known to be well (LKW)	38	38	36
Quality of In-hospital Stroke Care			
Early phase			
% of stroke patients who have brain imaging performed within 25 minutes of hospital arrival	67	71	69
% of ischemic stroke patients whose disease severity was assessed and documented using NIH stroke scale score	87	87	87
% of acute stroke patients who had dysphagia screening	87	87	86
% of acute stroke patients who received venous thromboembolism prophylaxis the day of or the day after admission	96	95	93
% of acute ischemic stroke patients who arrived at the hospital within 2 hours from time LKW and received IV alteplase within 3 hours of time LKW	90	95	92
% of eligible of acute ischemic stroke patients who received IV alteplase within 60 minutes of hospital arrival	85	87	85
% of ischemic stroke patients who received antithrombotic medication by the end of hospital day two	98	98	98
Later and at discharge			
% of ischemic stroke patients with medical history of smoking who received smoking cessation advice or counseling	99	99	99
% of ischemic stroke patients who were prescribed antithrombotic at discharge	99	99	99
% of ischemic stroke patients with atrial fibrillation who received anticoagulation therapy	97	97	96
% of eligible ischemic stroke patients who were discharged on statin medication	98	99	99
% of stroke patients who were assessed for rehabilitation services	99	99	99
Patient Education			
% of patients and/or caregiver that received educational materials during the hospital stay addressing all stroke education areas ¹	97	97	97
Aggregate			
% of patients with defect-free ² in-hospital care	79	79	78
Total number of patients	19,268	19,803	21,031

Abbreviation: **ED** – Emergency Department; **EMS** – Emergency Medical Service; **GCASR** – Georgia Coverdell Acute Stroke Registry; **LKW** – Last Known Well; **NIH** – National Institute of Health; **TIA** – Transient Ischemic Attack


¹Stroke education areas include activation of EMS, follow-up after discharge, medication adherence, risk factors, and sign and symptoms of stroke.

²Defect-free care is defined as the delivery of care meeting all quality indicators for which a patient is eligible.

References

1. Centers for Disease Control and Prevention, National Center for Health Statistics. Underlying Cause of Death 1999-2017 on CDC WONDER Online Database, released December, 2018. Data are from the Multiple Cause of Death Files, 1999-2016, as compiled from data provided by the 57 vital statistics jurisdictions through the Vital Statistics Cooperative Program. Accessed at <http://wonder.cdc.gov/ucd-icd10.html> on January 8, 2019.
2. Centers for Disease Control and Prevention (CDC). Prevalence and most common causes of disability among adults: United States, 2005. *Morbidity and Mortality Weekly Report*, 2009; 58:421–426. Available at <http://www.cdc.gov/mmwr/preview/mmwrhtml/mm5816a2.htm>
3. The National Institute of Neurological Disorders and Stroke rt-PA Stroke Study Group. Tissue plasminogen activator for acute ischemic stroke. *The New England Journal of Medicine*, 1995; 333:1581-1588. Available at: <http://www.nejm.org/doi/full/10.1056/NEJM199512143332401#t=article> doi: 10.1056/NEJM199512143332401
4. Benjamin EJ, Virani SS, Callaway CW, Chamberlain AM, Chang AR, Cheng S, et al; on behalf of the American Heart Association Council on Epidemiology and Prevention Statistics Committee and Stroke Statistics Subcommittee. Heart disease and stroke statistics—2017 update: a report from the American Heart Association. *Circulation*, 2018; 137(12):e67-e492. doi: 10.1161/CIR.0000000000000558.
5. Liao Y, Greenlund KJ, Croft JB, et al. Factors Explaining Excess Stroke Prevalence in the US Stroke Belt. *Stroke*, 2009; 40:3336-3341. Available at: <http://stroke.ahajournals.org/content/40/10/3336.full> (<https://doi.org/10.1161/STROKEAHA.109.561688>)
6. Georgia Behavioral Risk Factor Surveillance System Data 2017-2018. Chronic Disease, Healthy Behaviors, and Injury Epidemiology, Georgia Department of Public Health. For more information: <http://dph.georgia.gov/georgia-behavioral-risk-factor-surveillance-system-brfss>
7. Georgia Coverdell-Murphy Act. SB 549, Section 31-11-116. 14 May 2008, Official Code of Georgia Annotated, 2008. Available at: <http://www.legis.ga.gov/Legislation/20072008/85749.pdf>
8. Amendment to Georgia Coverdell-Murphy Act. House Bill 853, 26 April 2016. Accessed on December 15, 2017 at: <http://www.legis.ga.gov/Legislation/en-US/display/20152016/HB/853>
9. Ido MS, Bayakly R, Frankel M, Lyn R, Okosun IS. Administrative data linkage to evaluate a quality improvement program in acute stroke care, Georgia, 2006-2009. *Preventing Chronic Disease*, 2015; 12:E05. doi: 10.5888/pcd12.140238.
10. Ido MS, Okosun IS, Bayakly R, Clarkson L, Lugtu J, Floyd S, et al. Door to intravenous tissue plasminogen activator time and hospital length of stay in acute ischemic stroke patients, Georgia, 2007-2013. *Journal of Stroke and Cerebrovascular Disease*, 2016; 25(4):866-71. doi: 10.1016/j.jstrokecerebrovasdis.2015.12.025.
11. Ralph L. Sacco R, Emelia J. Benjamin EJ, Joseph P. Broderick JP, Mark Dyken M, J. Donald Easton JD, William M. Feinberg WM, et. Al. Risk Factors. *Stroke*, 1997; 28:1507-1517. Available at <http://stroke.ahajournals.org/content/28/7/1507.full> (<https://doi.org/10.1161/01.STR.28.7.1507>)
12. Population Projections, United States, 2004–2030, by state, age and sex, on CDC WONDER Online Database, September 2005. Accessed at <http://wonder.cdc.gov/population-projections.html> on December 15, 2018.

Map 1. Acute Care Hospitals participating in the Georgia Coverdell Acute Stroke Registry, November 2019



Definitions:

Alteplase: FDA-approved clot-busting drug for stroke. This drug can reduce disability by 30 percent in stroke sufferers if given to eligible patients within 3 hours of symptom onset.

Anticoagulation and Antithrombotic Medications: Medications that reduce blood clotting.

Atrial fibrillation: A disorder resulting in an irregular and often rapid heart rate. It predisposes to blood clotting and increases the risk of stroke, coronary heart disease and other heart-related complications.

Door-to-Needle Time: Time elapsed from when an eligible stroke patient arrives at the hospital to when Alteplase is administered. Eligible patients must receive Alteplase within 3 hours of symptom onset.

Dysphagia Screening: Screening for difficulty in swallowing. This identifies patients who need targeted treatment to improve their ability to swallow, so they do not aspirate or take fluid into the lungs. Aspiration of fluid can lead to pneumonia.

Hemorrhagic Stroke: A stroke caused by a blood vessel rupturing and bleeding in the brain. Hemorrhagic strokes are often fatal.

Ischemic Stroke: A stroke caused by a clot or blockage in a blood vessel supplying blood to the brain. The majority of strokes in Georgia are ischemic.

Statin medications: High cholesterol is a risk factor for stroke and statins lower blood cholesterol level.

Venous Thrombosis: When a blood clot forms in a vein, usually in the leg. If the clot breaks off, it can cause serious complications and even death.

Know the Signs and Symptoms of Heart Attack and Stroke

Heart attack and stroke are life-threatening emergencies. Call 911 if you experience these symptoms.

Signs of Heart Attack

- Chest discomfort. Most heart attacks involve discomfort in the center of the chest that lasts more than a few minutes, or that goes away and comes back. It can feel like uncomfortable pressure, squeezing, fullness, or pain.
- Discomfort in other areas of the upper body. Symptoms can include pain or discomfort in one or both arms, the back, neck, jaw, or stomach.
- Shortness of breath. This feeling often accompanies chest discomfort, but it can occur before the chest discomfort.
- Other symptoms may include nausea, lightheadedness, or breaking out in a cold sweat.

Signs of Stroke

- Sudden numbness or weakness of the face, arm, or leg, especially on one side of the body.
- Sudden confusion, trouble speaking or understanding.
- Sudden trouble seeing in one or both eyes
- Sudden trouble walking, dizziness, loss of balance or coordination.
- Sudden, severe headache with no known cause.