

Updated 4/23/2021

Week 15 (April 11 — April 17, 2021)

Snapshot of Influenza Activity During Week 15:

- Outpatient Illness Surveillance (ILINet): The proportion of outpatient visits for ILI was 1.9%, which is BELOW the regional baseline of 3.1%
- Activity Indicator Map: MINIMAL
- Influenza-associated Deaths: 0 deaths
- Metro Area Hospitalizations: 0 hospitalizations
- Influenza Outbreaks: 0 outbreaks
- **Viral Surveillance:** The percent of specimens testing positive for influenza by clinical laboratories was 0.2%

Summary of Select Influenza Surveillance Measures

	Week 15	Cumulative Data since September 27, 2020 (Week 40)*
No. of Influenza- associated Deaths	0	2
No. of Metro Area Influenza Hospitalizations	0	36
No. of Influenza Out- breaks	0	0

^{*}Cumulative data may include updated numbers from previous weeks.

Contact: Audrey.Kunkes@dph.ga.gov or (404) 463-4625




Outpatient Illness Surveillance


In Georgia during week 15, 1.9% of patient visits reported through the U.S. Outpatient Influenza-like Illness Surveillance Network (ILINet) were due to influenza-like illness (ILI). The percentage is below the regional baseline of 3.1%. (ILI is defined as fever (temperature of 100°F [37.8°C] or greater) and cough and/or sore throat.)

A total of 92 sentinel providers reported data for week 15.

The regional baseline is formulated by averaging ILI percentage during weeks of endemic activity determined by laboratory results for influenza. HHS Region 4 (AL, FL, GA, KY, MS, NC, SC, and TN) (Baseline: 3.1%).

Note: When drawing conclusions from the data please keep in mind that the influenza-like illness syndrome may be picking up COVID-19 illness.






Summary of ILI, by Age, Reported to ILINet, Weekly Georgia, 2020-2021 Influenza

Age Group in Years	No. of ILI Visits (Week 15)	Cumulative Data since September 27, 2020 (Week 40)
0-4	525	7,676
5-24	323	9,688
25-49	359	12,637
50-64	128	5,735
65+	82	4,081
Total	1,417	39,816

ILI Activity Indicator

ILI Activity Levels measure ILI activity each week. Activity levels are based on the percent of outpatient visits in Georgia due to ILI compared to the 3 year average of ILI visits during weeks with little or no influenza virus circulation.

During week 15, the activity level in Georgia was MINIMAL = 1



FLUVIEW interactive



A Weekly Influenza Surveillance Report Prepared by the Influenza Division

Influenza-Like Illness (ILI) Activity Level Indicator Determined by Data Reported to ILINet

2020-21 Influenza Season Week 15 ending Apr 17, 2021



Virgin Islands
*This map uses the proportion of outpatient visits to healthcare providers for influenza-like illness to measure the ILI activity level within a state. It does not, however, measure the extent of geographic spread of flu within a state. Therefore, outbreaks occurring in a single city could cause the state to display high activity levels.

*Data collected in ILINet may disproportionately represent certain populations within a state, and therefore may not accurately depict the full picture of influenza activity for the whole state.

*Data displayed in this map are based on data collected in ILINet, whereas the State and Territorial flu activity map are based on reports from state and territorial epidemiologists. The data presented in this map is preliminary and may change as more data is received.

*Differences in the data presented by CDC and state health departments likely represent differing levels of data completeness with data presented by the state likely being the more complete.

*For the data download you can use Activity Level for the number and Activity Level Label for the text description

*This graphic notice means that you are leaving an HHS Web site.

For more information, please see CDC's Exit Notification and Disclaimer policy.

For more information on the methodology, please visit Outpatient Illness Surveillance methods section.


Influenza-Associated Mortality

Influenza-associated deaths (in all ages) are reportable by law in the state of Georgia. To be confirmed as a as influenza-associated death, the person must have a clinically compatible illness, a positive influenza test, no documented recovery between the illness and death.

There were 0 influenza-associated death confirmed for week 15 in Georgia.

A total of 2 influenza-associated deaths have been confirmed for the 2020-2021 season.

Number of Laboratory Confirmed Influenza Deaths by Week of Death: Georgia Summary, 2020-2021 Influenza Season



Summary of Influenza-associated Deaths, by Age, Georgia, 2020-2021 Influenza Season


Age Group in Years	No. of Flu Deaths (Data Cumulative since Week 40)
0-4	0
5-17	0
18-49	0
50-64	1
65+	1
Total	2

Influenza-Associated Hospitalizations

The Influenza Hospitalization Surveillance Network (FluSurv-Net) reports laboratory confirmed influenza hospitalizations in the eight county metro Atlanta area (Fulton, DeKalb, Clayton, Cobb, Douglas, Gwinnett, Rockdale, and Newton) for the 2020-2021 influenza season.

There were 0 laboratory confirmed influenza hospitalizations confirmed for week 15.

A total of 36 laboratory confirmed influenza hospitalizations have been reported for the 2020-2021 season.



Summary of Influenza Hospitalizations, by Age, Georgia, 2020-2021 Influenza Season

Age Group in Years	No. of Flu Hospitalization (Cumulative Data since Week 40)	Hospitalization Rate (per 100,000 population)
0-4	1	0.37
5-17	0	0.00
18-49	6	0.32
50-64	14	1.86
65+	15	3.30
Total	36	0.88


Influenza Outbreaks

Influenza outbreaks are reportable by law in the state of Georgia.

0 influenza outbreaks were reported for week 15.

A total of 0 laboratory confirmed influenza outbreaks have been reported in Georgia for the 2020-2021 season.


Number of Laboratory Confirmed Influenza Outbreaks by Week Reported to Public Health, Weekly Georgia, Summary, 2020-2021 Influenza Season



Virologic Surveillance

The National Respiratory and Enteric Virus Surveillance System (NREVSS) and World Health Organization (WHO) collaborating laboratories (a combination of clinical and public health laboratories) report the total number of respiratory specimens tested for influenza and the number of positive for influenza, by virus type. Public Health Laboratories provide data about influenza virus subtypes and lineages (next page).


During week 15, clinical laboratories in Georgia reported testing 489 specimens, of which 0.2% were positive for influenza.



Summary of Influenza Tests from Clinical Laboratories, Georgia, 2020-2021 Influenza Season

	Week 15	Cumulative Data Since Week 40
No. of specimens tested	489	21,100
No. of positive specimens	1	132
Influenza A	1	50
Influenza B	0	82

During week 15, public health laboratories in Georgia reported testing 24 specimens for influenza, none were positive for influenza.




Summary of Influenza Tests from Public Health Laboratories, Georgia, 2020-2021 Influenza Season

	Cumulative Data Since Week 40
No. of specimens tested	4,183
No. of Positive Specimens	0
Influenza A (subtype not performed)	1
A(H1N1)pmd09	0
Н3	0
Influenza B (lineage not per- formed)	0
Yamagata lineage	0
Victoria lineage	0

Respiratory Syncytial Virus Infection (RSV) Surveillance

Data from NREVSS are also analyzed to measure the RSV seasonality. Antigen and polymerase chain reaction (PCR) tests are analyzed separately to determine the start and end of RSV season. Season onset is defined as the first week of two consecutive weeks when the percent positive of ALL laboratory confirmed tests are greater than or equal a certain threshold. The end is defined as the first week of two consecutive weeks when the percent positive of ALL laboratory confirmed tests are less than a certain threshold. For antigen-based testing, the threshold is 10% and for PCR the threshold is 3%.

During week 15, clinical laboratories in Georgia reported testing 18 specimens with antigen testing methods, 11.1% were positive for RSV.



During week 15, clinical laboratories in Georgia reported testing 958 specimens with PCR testing methods, 6.2% were positive for RSV.

