## 2006 Georgia Data Summary:

# **CARDIOVASCULAR DISEASE**



Cardiovascular disease is the leading cause of death in Georgia. However, there are many things you can do to reduce your risk and live a healthier life.

## CARDIOVASCULAR DISEASE MORBIDITY, MORTALITY, AND COSTS


#### **DEFINITION**

 Cardiovascular disease (CVD) includes all diseases of the heart and blood vessels, including ischemic heart disease, stroke, congestive heart failure, hypertensive disease, and atherosclerosis.

## **DEATHS**

- CVD accounted for one third (34%) of deaths in Georgia, with 22,378 CVD deaths in 2004.
- In 2004, Georgia's CVD death rate was 14% higher than the national rate.
- CVD death rates were 1.4 times higher for men than women in 2004.
- CVD death rates were 1.3 times higher for blacks than whites in 2004.

Cardiovascular Disease Death Rates by Race and Sex, Georgia, 2004




- CVD death rate increases with age but 1 in 4 persons who died from CVD in 2004 was less than 65 years of age.
- Black males are at particularly high risk for premature death from CVD. Almost 1 in 2 black males who died from CVD in 2004 was less than 65 years of age.

<sup>1</sup>Derived from national estimates published by the American Heart Association. *Heart Disease and Stroke Statistics - 2004 Update*. Dallas, TX: American Heart Association; 2004. <sup>2</sup>Hospital charges from acute care non-federal hospitals

## **DEATHS BY COUNTY**

Cardiovascular Disease Death Rates by County, Georgia, 2000-2004



## **ESTIMATED COSTS**

 The cost of CVD in Georgia in 2004 is estimated at \$9.6 billion,<sup>1</sup> which includes direct health care costs and lost productivity from morbidity and mortality (indirect costs).

## **HOSPITALIZATIONS**

- Approximately 145,000 hospitalizations occurred among Georgia residents due to CVD in 2004.
- The average charge per CVD hospitalization was \$26,200 in 2004.<sup>2</sup>
- Total hospital charges for CVD increased by over \$1.3 billion between 2001 and 2004, from \$2.5 billion to \$3.8 billion.
- The average hospitalization length of stay for CVD was 4 days in 2004.

## **HEART DISEASE AND STROKE RISK FACTORS**

## **SMOKING**

- 3,193 deaths due to CVD were attributable to smoking among adults ages 35 or older in 2004.
- Quitting smoking greatly reduces the risk for heart disease and stroke.
- In Georgia, 22% of adults reported smoking cigarettes in 2005.

## LACK OF PHYSICAL ACTIVITY

- Lack of physical activity increases the risk for heart disease and stroke.
- Regular physical activity helps to maintain proper body weight and improves general health.
- In 2005, 58% of adults in Georgia did not meet the Centers for Disease Control's recommendations for physical activity.

## POOR EATING HABITS

- Poor eating habits such as a high-fat diet can lead to obesity, ischemic heart disease, stroke and other chronic diseases.
- Eating five or more servings of fruits and vegetables a day can help prevent heart disease, cancer, and other chronic conditions.
- The majority (77%) of adults in Georgia did not consume the recommended five or more servings of fruits and vegetables per day in 2005.

## **OBESITY**

 Obesity, which is present in 27% of adults in Georgia, increases the risk of cardiovascular disease.

## **HIGH BLOOD PRESSURE**


- High blood pressure can by controlled through lifestyle changes such as physical activity and healthy diet and, if necessary, medications.
- The percentage of adults in Georgia who have been told that they have high blood pressure increased from 20% in 1995 to 27% in 2005.

## **HIGH CHOLESTEROL**

- Many people can control high cholesterol by modifying their diet. For those who cannot, medication can lower blood cholesterol levels.
- The percentage of adults in Georgia who have been told that they have high cholesterol increased from 22% in 1995 to 32% in 2005.

## **DIABETES**

- Diabetes can cause atherosclerosis, also known as hardening of the arteries
- Diabetes increases the risk for CVD death.
- The percentage of adults who report having diabetes increased from 4% in 1995 to 8% in 2005.



- (1) The percentage of Georgians who reported having been told they had high blood pressure
- (2) Of persons who had their blood cholesterol level checked, the percentage told that they have high cholesterol

Data Source: Georgia Behavioral Risk Factor Surveillance System, 2005

Date updated: August 2006 Publication number: DPH06.134HW

Visit http://www.health.state.ga.us/epi/cdiee/cardio.asp for more information about cardiovascular disease in Georgia.